

# Guidelines on Non-muscle-invasive **Bladder Cancer** (TaT1 and CIS)

M. Babjuk (chair), M. Burger, R. Zigeuner, S. Shariat,  
B. Van Rhijn, E. Comp erat, R. Sylvester, E. Kaasinen,  
A. Böhle, J. Palou, M. Roupr t

# TABLE OF CONTENTS

# PAGE

1.	BACKGROUND	4
1.1	Introduction	4
1.2	Methodology	4
1.2.1	Data Identification	4
1.2.2	Level of evidence and grade of recommendation	4
1.3	Publication history	5
1.3.1	Summary of changes	5
1.4	Potential conflict of interest statement	5
1.5	References	5
2.	EPIDEMIOLOGY	7
3.	RISK FACTORS	7
4.	CLASSIFICATION	7
4.1	Definition of non-muscle-invasive bladder cancer	7
4.2	Tumour, Node, Metastasis Classification (TNM)	8
4.3	Histological grading of non-muscle-invasive bladder urothelial carcinomas	8
4.4	Inter- and intraobserver variability in staging and grading	9
4.5	Further promising pathological parameters	10
4.6	Specific character of CIS and its clinical classification	10
5.	DIAGNOSIS	10
5.1	Patient history	10
5.2	Symptoms	10
5.3	Physical examination	10
5.4	Imaging	10
5.4.1	Intravenous urography and computed tomography	10
5.4.2	Ultrasonography	11
5.5	Urinary cytology	11
5.6	Urinary molecular marker tests	11
5.7	Practical application of urinary cytology and markers	12
5.8	Cystoscopy	12
5.9	Transurethral resection of Ta, T1 bladder tumours	13
5.10	Office-based fulguration	14
5.11	Bladder- and prostatic urethral biopsies	14
5.12	New TURB techniques	14
5.12.1	New resection techniques	14
5.12.2	New methods of tumour visualisation	14
5.12.2.1	Photodynamic diagnosis (fluorescence cystoscopy)	14
5.12.2.2	Narrow band imaging	15
5.13	Second resection	15
5.14	Pathological report	15
5.15	Guidelines for primary assessment of non-muscle-invasive bladder cancers	16
6.	PREDICTING DISEASE RECURRENCE AND PROGRESSION	17
6.1	Ta,T1 tumours	17
6.2	Carcinoma <i>in situ</i>	18
6.3	Recommendation for patients' stratification in risk groups	19
6.3.1	Recommendations for stratification of NMIBC	19
7.	ADJUVANT TREATMENT	19
7.1	Intravesical chemotherapy	19
7.1.1	One, immediate, postoperative intravesical instillation of chemotherapy	19
7.1.2	Additional adjuvant intravesical chemotherapy instillations	20
7.1.3	Options for improving efficacy of intravesical chemotherapy	20
7.2	Intravesical Bacillus Calmette-Guérin (BCG) immunotherapy	20
7.2.1	Efficacy of BCG	20

7.2.2	Optimal BCG schedule	21
7.2.3	BCG toxicity	21
7.2.4	Optimal dose of BCG	23
7.2.5	BCG strain	23
7.2.6	Indications for BCG	23
7.3	Specific aspects of treatment of CIS	23
7.3.1	Treatment strategy	23
7.3.2	Cohort studies	23
7.3.3	Prospective randomised trials	23
7.3.4	Treatment of extravesical CIS	24
7.4	Treatment of failure of intravesical therapy	24
7.4.1	Failure of intravesical chemotherapy	24
7.4.2	Recurrence and failure after intravesical BCG immunotherapy	24
7.4.3	Treatment of BCG failure and recurrences after BCG	24
7.5	Recommendations for adjuvant therapy in Ta, T1 tumours and for therapy of CIS	25
8.	RADICAL CYSTECTOMY FOR NON-MUSCLE-INVASIVE BLADDER CANCER	25
9.	FOLLOW-UP OF PATIENTS WITH NON-MUSCLE-INVASIVE BLADDER TUMOURS	27
10.	REFERENCES	28
11.	ABBREVIATIONS USED IN THE TEXT	42

# 1. BACKGROUND

## 1.1 Introduction

This overview represents the updated European Association of Urology (EAU) guidelines for Non-muscle-invasive Bladder Cancer (CIS, Ta, T1). The information presented is limited to urothelial carcinoma, if not specified otherwise. Aim is to provide practical guidance on the clinical management of non-muscle-invasive bladder cancer with a focus on clinical presentation and recommendations.

The EAU Guidelines Panel on Non-muscle-invasive Bladder Cancer consists of an international multidisciplinary group of clinicians, including a pathologist and a statistician.

It must be emphasised that clinical guidelines present the best evidence available but following the recommendations will not necessarily result in the best outcome. Guidelines can never replace clinical expertise when making treatment decisions for individual patients - also taking personal values and preferences and individual circumstances of patients into account.

Separate EAU guidelines documents are available addressing upper urinary tract tumours (1), muscle-invasive bladder cancer (2), and primary urethral carcinomas (3).

## 1.2 Methodology

### 1.2.1 Data Identification

The systematic literature search for each section of the Non-muscle-invasive Bladder Cancer Guidelines was performed by the panel members. For identification of original and review articles, Medline, Web of Science, and Embase databases were used. For the current update, all articles published between 2010 and 2012 on non-muscle-invasive bladder cancer were considered. Focus of the searches was identification of all level 1 scientific papers (randomised controlled trials (RCTs), systematic reviews (SRs), and meta-analyses of RCTs) in accordance with the EAU guidelines methodology.

### 1.2.2 Level of evidence and grade of recommendation

References in the text have been assessed according to their level of scientific evidence (LE), and guideline recommendations have been graded follow the listings in Tables 1 and 2, based on the Oxford Centre for Evidence-based Medicine Levels of Evidence (4). Grading aims to provide transparency between the underlying evidence and the recommendation given.

**Table 1: Level of evidence\***

Level	Type of evidence
1a	Evidence obtained from meta-analysis of randomised trials.
1b	Evidence obtained from at least one randomised trial.
2a	Evidence obtained from one well-designed controlled study without randomisation.
2b	Evidence obtained from at least one other type of well-designed quasi-experimental study.
3	Evidence obtained from well-designed non-experimental studies, such as comparative studies, correlation studies and case reports.
4	Evidence obtained from expert committee reports or opinions or clinical experience of respected authorities.

\*Modified from Sackett et al. (4).

It should be noted that when recommendations are graded, the link between the LE and grade of recommendation (GR) is not directly linear. Availability of RCTs may not necessarily translate into a grade A recommendation where there are methodological limitations or disparity in published results.

Alternatively, absence of high level of evidence does not necessarily preclude a grade A recommendation, if there is overwhelming clinical experience and consensus. There may be exceptional situations where corroborating studies cannot be performed, perhaps for ethical or other reasons and in this case unequivocal recommendations are considered helpful. Whenever this occurs, it is indicated in the text as “upgraded based on panel consensus”. The quality of the underlying scientific evidence - although a very important factor - has to be balanced against benefits and burdens, values and preferences, and costs when a grade is assigned (5-7).

The EAU Guidelines Office does not perform structured cost assessments, nor can they address local/national preferences in a systematic fashion. But whenever these data are available, the expert panel will include the information.

**Table 2: Grade of recommendation\***

Grade	Nature of recommendations
A	Based on clinical studies of good quality and consistency that addressed the specific recommendations, including at least one randomised trial.
B	Based on well-conducted clinical studies, but without randomised clinical trials.
C	Made despite the absence of directly applicable clinical studies of good quality.

\*Modified from Sackett et al. (4).

### 1.3 Publication history

The first European Association of Urology (EAU) Guidelines on Bladder Cancer were published in 2000 (8). In 2004 it was decided to develop separate guidelines for muscle-invasive and infiltrative bladder cancer and a separate scientific publication on upper urinary tract tumours was presented (9), which was updated and has been included in the EAU Guidelines compilation print since 2011 (10). The complete updates of guidelines for non-muscle-invasive bladder cancer were prepared in 2006, 2008 and 2011 (11-14). Since 2011 the EAU Guidelines on TaT1 tumours and CIS were integrated in one guidelines document (14).

Several scientific summaries have been published in the EAU scientific journal, *European Urology* (9,15-19). A quick reference document (pocket guidelines) is available presenting the main findings of the Non-muscle-invasive Bladder Cancer Guidelines. This document follows the updating cycle of the underlying large texts.

All material can be viewed and downloaded for personal use at the EAU website. The EAU website also includes a selection of translations and republications produced by national urological associations: <http://www.uroweb.org/guidelines/online-guidelines/>.

This document was peer-reviewed prior to publication.

#### 1.3.1 Summary of changes

For all chapters the literature has been assessed.

Chapter 4 - Classification: a clear definition of non-muscle-invasive bladder cancer is presented. Since appropriate classification and grading directly influences treatment decisions, additional information on pathological parameters has been added.

Chapter 5 - Diagnosis: an illustration on bladder diagram to facilitate the description of cystoscopy finding has been added. The new data on endoscopic diagnosis and pathological evaluation of the tissue included in this section resulted in a number of changes in the recommendations.

Chapter 6 - Predicting disease recurrence and progression: the new stratification of patients into 3 risk groups facilitating treatment recommendation is presented

Chapter 7 - Adjuvant treatment: updated information on intravesical chemo- and immunotherapy is provided. The definition and stratification of BCG toxicity and side-effects is provided in an overview table. The definition of BCG failures has been specified.

Chapter 8: Radical cystectomy for NMIBC: the indication criteria were updated.

### 1.4 Potential conflict of interest statement

The expert panel have submitted potential conflict of interest statements which can be viewed on the EAU website: <http://www.uroweb.org/guidelines/>.

### 1.5 References

1. Rouprêt M, Babjuk M, Compérat E, et al. EAU Guidelines on Urothelial Carcinomas of the Upper Urinary Tract. In: EAU Guidelines, edn. presented at the EAU Annual Congress Milan 2013. ISBN 978-90-79754-71-7.
2. Witjes JA, Compérat E, Cowan NC, et al. EAU Guidelines on Muscle-invasive and Metastatic Bladder Cancer. In: EAU Guidelines, edn. presented at the EAU Annual Congress Milan 2013. ISBN 978-90-79754-71-7.
3. Gakis G, Witjes JA, Compérat E, et al. Guidelines on Primary Urethral Carcinoma. Edition presented at the EAU Annual Congress Milan 2013. ISBN: 978-90-79754-71-7.

4. Oxford Centre for Evidence-based Medicine Levels of Evidence (May 2009). Produced by Bob Phillips, Chris Ball, Dave Sackett, Doug Badenoch, Sharon Straus, Brian Haynes, Martin Dawes since November 1998. Updated by Jeremy Howick March 2009.  
<http://www.cebm.net/index.aspx?o=1025> (Access date January 2013)
5. Atkins D, Best D, Briss PA, et al; GRADE Working Group. Grading quality of evidence and strength of recommendations. *BMJ* 2004 Jun 19;328(7454):1490.  
<http://www.ncbi.nlm.nih.gov/pubmed/15205295>
6. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336(7650):924-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/18436948>
7. Guyatt GH, Oxman AD, Kunz R, et al; GRADE Working Group. Going from evidence to recommendations. *BMJ* 2008 May 10;336(7652):1049-51.  
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2376019/>
8. Oosterlinck W, Jakse G, Malmström P-U, et al. EAU Guidelines on Bladder Cancer. In EAU Guidelines, edn. presented at the EAU Annual Congress Brussels, 2000.
9. Oosterlinck A, Van der Meijden A, Sylvester R et al. EAU Guidelines on TaT1 (non-muscle-invasive) Bladder Cancer. In: EAU Guidelines, edn. presented at the EAU Annual Congress Paris, 2006. ISBN 90-70244-37-3.
10. Babjuk M, Oosterlinck W, Sylvester R, et al EAU Guidelines on TaT1 (Non-muscle-invasive) Bladder Cancer. In: EAU Guidelines edn. presented at the EAU Annual Congress Milan, 2008. ISBN 978-90-70244-91-0.
11. Babjuk M, Oosterlinck W, Sylvester R, et al EAU Guidelines on TaT1 (non-muscle invasive) Bladder Cancer. In EAU Guidelines, edn. presented at the EAU Annual Congress Stockholm 2009. ISBN 978-90-79754-09-0.
12. Babjuk M, Oosterlinck W, Sylvester R, et al. EAU Guidelines on Non-muscle-invasive Bladder Cancer (TaT1 and CIS). In: EAU Guidelines, edn. presented at the EAU Annual Congress Congress Vienna, 2011. ISBN 978-90-79754-83-0.
13. Oosterlinck W, Solsona E, van der Meijden AP, Sylvester R, Böhle A, Rintala E, Lobel B; European Association of Urology. EAU guidelines on diagnosis and treatment of upper urinary tract transitional cell carcinoma. *Eur Urol.* 2004 Aug;46(2):147-54.  
<http://www.ncbi.nlm.nih.gov/pubmed/15245806>
14. Rouprêt M, Zigeuner R, Palou J, et al. Guidelines on Upper Urinary Tract Urothelial Cell Carcinomas. In: EAU Guidelines, edn presented at the EAU Annual Congress Vienna, 2011. ISBN 978-90-79754-83-0.
15. Oosterlinck W, Lobel B, Jakse G, Malmström PU, Stöckle M, Sternberg C; European Association of Urology (EAU) Working Group on Oncological Urology. Guidelines on bladder cancer. *Eur Urol* 2002 Feb;41(2):105-12.  
<http://www.ncbi.nlm.nih.gov/pubmed/12074395>
16. Van der Meijden AP, Sylvester R, Oosterlinck W, et al; for the EAU Working Party on Non Muscle Invasive Bladder Cancer. EAU guidelines on the diagnosis and treatment of urothelial carcinoma in situ. *Eur Urol.* 2005 Sep;48(3):363-71.  
<http://www.ncbi.nlm.nih.gov/pubmed/15994003>
17. Babjuk M, Oosterlinck W, Sylvester R, et al; European Association of Urology (EAU). EAU guidelines on non-muscle-invasive urothelial carcinoma of the bladder. *Eur Urol* 2008 Aug;54(2):303-14.  
<http://www.ncbi.nlm.nih.gov/pubmed/18468779>
18. Rouprêt M, Zigeuner R, Palou J, et al. European guidelines for the diagnosis and management of upper urinary tract urothelial cell carcinomas: 2011 update. *Eur Urol* 2011 Apr;59(4):584-94.  
<http://www.ncbi.nlm.nih.gov/pubmed/21269756>
19. Babjuk M, Oosterlinck W, Sylvester R, et al; European Association of Urology (EAU). EAU guidelines on non-muscle-invasive urothelial carcinoma of the bladder, the 2011 update. *Eur Urol* 2011 Jun;59(6):997-1008.  
<http://www.ncbi.nlm.nih.gov/pubmed/21458150>

## 2. EPIDEMIOLOGY

Bladder Cancer (BC) is the most common malignancy of the urinary tract and the seventh most common cancer in men and the 17<sup>th</sup> in women. The worldwide age standardised incidence rate is 9 per 100,000 for men and 2 per 100,000 for women (2008 data) (1). In the European Union (EU) age standardised incidence rate is 27 per 100,000 for men and six per 100,000 for women (1).

Incidence varies between regions and countries; in Europe, the highest age standardised incidence rate has been reported in Spain (41.5 in men and 4.8 in women) and the lowest in Finland (18.1 in men and 4.3 in women) (1). The variations can partly be attributed to different methodology and quality of data collection, thus warranting care in the interpretation of results (2,3).

The world global age standardised mortality rate is 3 for men versus 1 per 100,000 for women. In the EU, age standardised mortality rate is 8 for men and 3 per 100,000 for women, respectively (1). In 2008 BC was the eighth most common cause of cancer-specific mortality in Europe (1).

The incidence of BC has decreased in some registries possibly reflecting decreased impact of causative agents, mainly smoking and occupational exposure (4). The mortality of BC has also decreased, possibly reflecting increased standard of care (5).

Approximately 75% of patients with BC present with a disease that is confined to the mucosa (stage Ta, CIS) or submucosa (stage T1). These categories are grouped as non-muscle-invasive bladder tumours. Non-muscle invasive BC (NMIBC) has a high prevalence due to low progression rates and long-term survival in many cases; patients with muscle-invasive BC (MIBC) are at higher risk of cancer-specific mortality (3). The prevalence of BC is among the highest of all urological malignancies (1).

## 3. RISK FACTORS

Increasing evidence suggests that genetic predisposition has a significant influence on bladder cancer incidence, especially via its impact on susceptibility to other risk factors (3,6). Tobacco smoking is the most important risk factor for BC, accounting for ~50% of cases (3,7) (LE: 3). Tobacco smoke contains aromatic amines and polycyclic aromatic hydrocarbons, which are renally excreted.

Occupational exposure to aromatic amines, polycyclic aromatic hydrocarbons and chlorinated hydrocarbons is the second most important risk factor for BC, accounting in the modern era for ~10% of all cases. Such occupational exposure occurs mainly in industrial branches processing paint, dye, metal and petroleum products (3,8-10) (LE: 3).

Although the significance of the amount of fluid intake is uncertain, chlorination of drinking water and subsequent levels of trihalomethanes is potentially carcinogenic, and exposure to arsenic in drinking water increases BC risk (3,11) (LE: 3). The relation between personal hair dye use and BC risk remains uncertain; increased risk has been suggested in users of permanent hair dyes with NAT2 slow acetylation phenotype (12,13).

The exposure to ionizing radiation is connected with increased risk of BC (LE: 3). It is suggested that cyclophosphamide and pioglitazone are weakly associated with BC risk (3). Schistosomiasis, a chronic endemic cystitis based on recurrent infection with a parasitic trematode, is a cause of bladder cancer, particularly squamous cell carcinoma (3) (LE: 3).

## 4. CLASSIFICATION

### 4.1 Definition of non-muscle-invasive bladder cancer

A papillary tumour confined to the mucosa is classified as stage Ta according to the Tumour, Node, Metastasis (TNM) classification system. Tumours that have invaded the lamina propria are classified as stage T1. Ta and T1 tumours can be removed by transurethral resection (TUR), and therefore they are grouped under the heading of NMIBC for therapeutic purposes. Also included under this heading are flat, high-grade tumours that are confined to the mucosa, and classified as CIS (Tis). However, molecular biology techniques and clinical experience have demonstrated the highly malignant potential of CIS and T1 lesions. Therefore, the terms NMIBC and superficial BC are suboptimal descriptions. The latter term should no longer be used. Whenever the terminology NMIBC is used in individual cases, the tumour stage and grade should be mentioned.

## 4.2 Tumour, Node, Metastasis Classification (TNM)

The 2002 TNM classification approved by the Union International Contre le Cancer (UICC) has been widely accepted. This version was updated in 2009, but it has no changes for bladder tumours (Table 3) (14).

**Table 3: 2009 TNM classification of urinary bladder cancer**

<b>T - Primary tumour</b>	
TX	Primary tumour cannot be assessed
T0	No evidence of primary tumour
Ta	Non-invasive papillary carcinoma
Tis	Carcinoma <i>in situ</i> : 'flat tumour'
T1	Tumour invades subepithelial connective tissue
T2	Tumour invades muscle
T2a	Tumour invades superficial muscle (inner half)
T2b	Tumour invades deep muscle (outer half)
T3	Tumour invades perivesical tissue:
T3a	Microscopically
T3b	Macroscopically (extravesical mass)
T4	Tumour invades any of the following: prostate, uterus, vagina, pelvic wall, abdominal wall
T4a	Tumour invades prostate, uterus or vagina
T4b	Tumour invades pelvic wall or abdominal wall
<b>N - Lymph nodes</b>	
NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in a single lymph node in the true pelvis (hypogastric, obturator, external iliac, or presacral)
N2	Metastasis in multiple lymph nodes in the true pelvis (hypogastric, obturator, external iliac, or presacral)
N3	Metastasis in common iliac lymph node(s)
<b>M - Distant metastasis</b>	
MX	Distant metastasis cannot be assessed
M0	No distant metastasis
M1	Distant metastasis

## 4.3 Histological grading of non-muscle-invasive bladder urothelial carcinomas

In 1998, a new classification of non-invasive urothelial tumours was proposed by the World Health Organization (WHO) and the International Society of Urological Pathology (ISUP) (1998 WHO/ISUP classification) and published by the WHO in 2004 (15,16) (Table 4). Its major contribution is a detailed histological description of the various grades, which uses specific cytological and architectural criteria. A website ([www.pathology.jhu.edu/bladder](http://www.pathology.jhu.edu/bladder)) that illustrates examples of the various grades has been developed to further improve accuracy in using the system.



**Table 4: WHO grading in 1973 and in 2004 (15,16)**


<p><b>1973 WHO grading</b>                  Urothelial papilloma                  Grade 1: well differentiated                  Grade 2: moderately differentiated                  Grade 3: poorly differentiated</p> <p><b>2004 WHO grading</b>                  Flat lesions                  Hyperplasia (flat lesion without atypia or papillary aspects)                  Reactive atypia (flat lesion with atypia)                  Atypia of unknown significance                  Urothelial dysplasia                  Urothelial CIS                  Papillary lesions                  Urothelial papilloma (completely benign lesion)                  Papillary urothelial neoplasm of low malignant potential                  Low-grade papillary urothelial carcinoma                  High-grade papillary urothelial carcinoma</p>
--

The 2004 WHO classification of the flat lesions includes urothelial hyperplasia, reactive urothelial atypia, atypia of unknown significance, dysplasia, and CIS. Among non-invasive papillary urothelial lesions, the 2004 WHO grading differentiates between papillary urothelial neoplasm of low malignant potential (PUNLMP) and low-grade and high-grade urothelial carcinomas.

Papillary urothelial neoplasms of low malignant potential (PUNLMPs) are defined as lesions that do not have cytological features of malignancy but show normal urothelial cells in a papillary configuration. Although they have a negligible risk for progression, they are not completely benign and still have a tendency to recur. The intermediate grade (Grade 2), which was the subject of controversy in the 1973 WHO classification, has been eliminated (17-19) (Figure 1). The published comparisons, however, have not clearly confirmed that the WHO 2004 classification has better reproducibility than the 1973 classification (20,21).

The prognostic value of both grading systems (WHO 1973 and 2004) has been confirmed. Attempts to demonstrate better prognostic value of one of the systems, however, have yielded controversial results (17-20,22-24). The majority of clinical trials published to date on TaT1 bladder tumours have been performed using the 1973 WHO classification, and therefore, the following guidelines are based on this version. Until the prognostic role of WHO 2004 is validated by more prospective trials, both classifications should be used.

**Figure 1: Stratification of tumours according to grade in the WHO 1973 and 2004 classifications (19)\***



**Histologic Spectrum of transitional cell carcinoma (urothelial carcinoma [UC] & spectrum)**

*\*1973 WHO Grade 1 carcinomas have been reassigned to PUNLMP and Low-grade carcinomas in 2004 WHO classification, and Grade 2 carcinomas to Low-grade and High-grade carcinomas. All 1973 WHO Grade 3 carcinomas have been reassigned to High-grade carcinomas. Reproduced with permission from Elsevier.*

**4.4 Inter- and intraobserver variability in staging and grading**

Despite well-defined criteria for the diagnosis of urothelial carcinoma, there is significant variability among pathologists for diagnosis of CIS, for which agreement is achieved in only 70-78% of cases (25,26) (LE: 2a). There is interobserver variability in classification of stage T1 versus Ta tumours and tumour grading in both 1997 and 2004 classifications. The general conformity in staging and grading is between 50 and 60% (20,25-29) (LE: 2a).

In difficult cases, an additional review by an experienced genitourinary pathologist is recommended.

#### **4.5 Further promising pathological parameters**

Some novel parameters based on pathological investigation of resected tissue have been evaluated and considered for subclassification and prognostic purposes.

In patients with T1 tumours, the depth and extent of invasion into the lamina propria (T1 substaging) can be evaluated. The prognostic value of this evaluation has been demonstrated by some retrospective cohort studies (30-33) (LE: 3). The presence of lymphovascular invasion has been reported as an unfavourable prognostic factor in T1 tumours (34) (LE: 3). It must be presented in pathological reports.

Detection of the micropapillary variant of urothelial carcinoma represents a poor prognostic factor even if it is non-muscle invasive at the time of diagnosis (35,36) (LE: 3). In micropapillary urothelial tumours with T1 invasion, some cases with distant metastases have been confirmed (35). Moreover, the risk of understaging in these tumours is substantial (37).

Rare cases of non-invasive squamous cell carcinoma in the bladder with poor prognosis have been described (38). Novel molecular markers, particularly FGFR3 mutation status, are promising but need further evaluation (17,33,39-41).

#### **4.6 Specific character of CIS and its clinical classification**

CIS is a flat, high-grade, non-invasive urothelial carcinoma. Macroscopically, CIS can be missed at cystoscopy or be considered as an inflammatory lesion if it is not biopsied. It is often multifocal and can occur not only in the bladder but also in the upper urinary tract, prostatic ducts, and prostatic urethra (42).

CIS is classified into one of four different clinical types (43):

- Primary: isolated CIS with no previous or concurrent papillary tumours and no previous CIS;
- Secondary: CIS detected during follow-up of patients with a previous tumour that was not CIS;
- Concurrent: CIS in the presence of any other urothelial tumour in the bladder;
- Recurrent: Repeat occurrence of isolated CIS after initial successful response to intravesical treatment.

## **5. DIAGNOSIS**

### **5.1 Patient history**

Patient history should be taken and recorded for all important information with possible connection to BC, including risk factors and history of suspect symptoms.

### **5.2 Symptoms**

Haematuria is the most common finding in NMIBC. Ta, T1 tumours do not cause bladder pain and rarely present with lower urinary tract symptoms (LUTS). In patients who do complain of these symptoms, particularly in those with irritative LUTS refractory to symptomatic treatment, CIS might be suspected.

### **5.3 Physical examination**

Physical examination does not reveal NMIBC.

### **5.4 Imaging**

#### **5.4.1 Intravenous urography and computed tomography**

Intravenous urography (IVU) is used to detect filling defects in the calyces, renal pelvis and ureters, and hydronephrosis, which can indicate the presence of a ureteral tumour. Large exophytic tumours may be seen as filling defects in the bladder. The necessity to perform routine IVU once a bladder tumour has been detected is questioned because of the low incidence of significant findings obtained with this method (44-46) (LE: 2a). The incidence of upper urinary tract tumours is low (1.8%), but increases to 7.5% in tumours located in the trigone (45) (LE: 2b). The risk of tumour recurrence in the upper urinary tract during follow-up increases in multiple and high-risk tumours (47) (LE: 2b).

In most centres, computed tomography (CT) urography is used as an alternative to conventional IVU (48). Especially in muscle-invasive tumours of the bladder and upper urinary tract tumours, CT urography gives more information than IVU does (including status of lymph nodes and neighbouring organs). However, CT urography has the disadvantage of higher radiation exposure compared to IVU.

#### 5.4.2 **Ultrasonography**

Ultrasonography (US) is often used as the initial tool to assess the urinary tract. This is not only because it avoids the use of contrast agents, but also because sensitive transducers have improved imaging of the upper urinary tract and bladder.

Transabdominal US permits characterisation of renal masses, detection of hydronephrosis, and visualisation of intraluminal masses in the bladder. It can be as accurate as IVU for diagnosis of upper urinary tract obstruction (44) (LE: 3). US is therefore a useful tool for detection of obstruction in patients with haematuria, however, it cannot exclude the presence of upper tract tumours.

CIS cannot be diagnosed with imaging methods (IVU, CT urography or US).

### 5.5 **Urinary cytology**

Examination of voided urine or bladder-washing specimens for exfoliated cancer cells has high sensitivity in high-grade tumours but low sensitivity in low-grade tumours. As a result of loss of cell cohesion in the epithelial lining of the bladder in CIS, there is a larger number of floating cells in the urine, as well as a high degree of anaplasia. The sensitivity of cytology for CIS detection is 28-100% (49) (LE: 2b). Cytology is thus useful when a high-grade malignancy or CIS is present. However, urinary cytology often is negative in the presence of low-grade cancer. Positive voided urinary cytology can indicate a urothelial tumour anywhere in the urinary tract, from the calyx to the ureters, bladder, and proximal urethra. Negative cytology, however, does not exclude the presence of a tumour in the urinary tract.

Cytological interpretation is user-dependent (50). Evaluation can be hampered by low cellular yield, urinary tract infections, stones, or intravesical instillations. In experienced hands however, the specificity exceeds 90% (51) (LE: 2b). Cytology should be performed on fresh urine with adequate fixation. Morning urine is not suitable because of the frequent presence of cytolysis.

### 5.6 **Urinary molecular marker tests**

There are specified general requirements for good bladder cancer markers (51):

- The test must be as technically simple as possible (preferably a point-of-care test, with readily available results, easy to perform, with a short learning curve);
- Low cost;
- Good reliability and reproducibility;
- For individual patient populations and clinical situations, the test should have a high specificity to avoid unnecessary workup because of false-positive results, and high sensitivity to avoid the risk of missing a tumour;
- For clinical settings, it is of utmost importance to detect high-risk urothelial cancer before it escapes curative treatment.

Driven by the low sensitivity of urine cytology, extensive laboratory research has developed numerous urinary tests for BC detection (51-57). Considering the frequency of cystoscopy for follow-up, markers for recurrent urothelial cancer would be especially useful.

Numerous reviews of urinary markers have appeared in recent years (51-53,55-65). None of these markers have been accepted as standard diagnostic or follow-up procedures in routine urology or in guidelines. Some urine tests that have been evaluated in several laboratories/centres and in studies with sufficient numbers of patients are listed in Table 5. Sensitivity and specificity should be used to compare studies on urine tests because they remain constant, whereas positive and negative predictive values vary between populations with different numbers of positive and negative events (54,57).

The following conclusions can be drawn about the existing tests. Sensitivity is usually higher at the cost of lower specificity than urine cytology (51-65) (LE: 3). Benign conditions and BCG influence many urinary marker tests (51-65) (LE: 3). Sensitivity and specificity of a urinary marker test depend on the clinical context of the patient (screening, primary detection, follow-up (high risk), and follow-up (low/intermediate risk)) (54-57) (LE: 3). For example, sensitivity of a given urinary marker is higher for detection of a primary lesion than a recurrent lesion (54) (LE: 3). Patient selection explains the wide range in performance of the markers listed in Table 5.

Unlike other urine tests, some false-positive results of UroVysion and microsatellite analysis can be attributed to occult disease and thus identify those patients who are more likely to experience subsequent recurrence. It might also be useful to predict response to intravesical therapy (66-68) (LE: 3). Microsatellite analysis is the most promising of the methods listed in Table 5 (69-71).

**Table 5: Summary of main urinary markers**

Markers (or test specifications)	Overall sensitivity (%)	Overall specificity (%)	Sensitivity for high-grade tumours (%)	Point-of-care test	Level of evidence (LE)
UroVysion	30-86	63-95	66-70	No	3
Microsatellite analysis	58 -92	73-100	90-92	No	1b
Immunocyt/uCyt +	52-100	63-75	62-92	No	3
Nuclear matrix protein 22	47-100	55-98	75-83	Yes	3
BTA stat	29-83	56-86	62-75	Yes	3
BTA TRAK	53-91	28-83	74-77	No	3
Cytokeratins	12-88	73-95	33-100	No	3

BTA = bladder tumour antigen.

### 5.7 Practical application of urinary cytology and markers

The following objectives of urinary cytology or molecular tests must be considered:

- Screening of the population at risk of BC.  
The application of haematuria dipstick, NMP22 or UroVysion in BC screening in high-risk populations has been reported (72,73). The low incidence of BC in the general population and the short lead time impair feasibility and cost-effectiveness (57,72-74). Routine application of screening is not recommended.
- Exploration of patients after haematuria or other symptoms suggestive of BC (primary detection).  
It is generally accepted that none of the tests can replace cystoscopy. However, urinary cytology or markers can be used as an adjunct to cystoscopy to detect invisible tumours, particularly CIS. In this setting, sensitivity for high-grade tumours and specificity are particularly important. Urinary cytology is highly specific but urinary markers lack this high specificity and are not recommended for primary detection. Future studies should explore the feasibility of urine markers preceding/replacing cystoscopy in patients with microscopic haematuria.
- Facilitate surveillance of NMIBC (54,59,75,76).
  - a. Follow-up of high-risk NMIBC:  
High-risk tumours should be detected early in follow-up, and the percentage of tumours missed should be as low as possible. Therefore, the best surveillance strategy for these patients will continue to include frequent cystoscopy and cytology. Specificity is more important than sensitivity in this subset of patients, because the urinary markers are used as an adjunct to cystoscopy. A urinary marker other than cytology is not recommended for high-risk NMIBC surveillance.
  - b. Follow-up low/intermediate risk NMIBC:  
To reduce the number of cystoscopy procedures, urinary markers should be able to detect recurrence before the tumours are large and numerous. The limitation of urinary cytology is its low sensitivity for low grade/risk recurrences. Several urinary markers are better but still do not detect half of the low-grade tumours that are detected by cystoscopy (54,57) (LE: 3).

According to current knowledge, no urinary marker can replace cystoscopy during follow-up or help to lower cystoscopic frequency in routine fashion. One prospective randomised study confirmed that knowledge of positive test results (microsatellite analysis) can improve the quality of follow-up cystoscopy (77) (LE: 1b). It supports the adjunctive role of a non-invasive urine test performed before follow-up cystoscopy. (77).


### 5.8 Cystoscopy

The diagnosis of papillary BC ultimately depends on cystoscopic examination of the bladder and histological evaluation of the resected tissue. CIS is diagnosed by a combination of cystoscopy, urine cytology, and histological evaluation of multiple bladder biopsies (78).

Cystoscopy is initially performed in the office. A flexible instrument with topical intraurethral anaesthetic lubricant instillation results in better compliance, especially in men (79). Careful inspection of the whole urothelial lining in the bladder should be performed to prevent missing the tumour.

If a bladder tumour has been visualised in earlier imaging studies, diagnostic cystoscopy can be omitted because the patient will undergo TUR (80).

A careful description of the findings is necessary. It should include the site, size, number, and appearance (papillary or solid) of the tumours, as well as a description of mucosal abnormalities. Use of a bladder diagram is recommended (Figure 2).



1 = Trigone	6 = Anterior wall
2 = Right ureteral orifice	7 = Posterior wall
3 = Left ureteral orifice	8 = Dome
4 = Right wall	9 = Neck
5 = Left wall	10 = Posterior urethra

**Figure 2: Bladder diagram**

### 5.9 Transurethral resection of Ta, T1 bladder tumours

The goal of the TURB in Ta, T1 BC is to make the correct diagnosis and remove all visible lesions. It is a crucial procedure in the diagnosis and treatment of BC.

TUR of the bladder (TURB) should be performed systematically as follows:

- Procedure is initiated with careful bimanual palpation under general or spinal anaesthesia;
- Insertion of the resectoscope, in men under visual guidance, with inspection of the whole urethra;
- Inspection of the whole urothelial lining of the bladder;
- Biopsy from prostatic urethra (if indicated, see lower);
- Cold-cup bladder biopsies (if indicated, see lower);
- Resection of the tumour.

The strategy of resection depends on the size of the lesion. Small tumours (< 1 cm) can be resected *en bloc*, which includes the entire tumour and part of the underlying bladder wall. Larger tumours should be resected separately in fractions, including the exophytic part of the tumour, the underlying bladder wall with the detrusor muscle, and the edges of the resection area. This approach provides good information about the vertical and horizontal extent of the tumour and helps to improve resection completeness (81) (LE: 3). Deep resection is not necessary in small, apparently low-grade lesions with a previous history of low-grade Ta (G1) tumour.

- In patients with palpable lesions before TURB, bimanual palpation should be repeated after resection
- The protocol is formulated, which must describe all previous steps of the procedure, as well as extent and completeness of resection
- An order form for pathological evaluation is prepared.

The specimens from different biopsies and resection fractions must be referred to the pathologist in separate containers and labelled separately, to enable him/her to make a correct diagnosis. Cauterisation should be avoided as much as possible during TURB to prevent tissue destruction.

Complete and correct TURB is essential to achieve a good prognosis (82). It has been confirmed that absence of detrusor muscle in the specimen is associated with a significantly higher risk of residual disease

and early recurrence (83) (LE: 2b).

Training in the methods of TURB should be included in teaching programmes. It has been shown that surgical experience can improve TURB results (84).

### 5.10 Office-based fulguration

In patients with a history of small, low-grade (WHO 2004 classification)/G1 (1973 classification) Ta tumours, fulguration of small papillary recurrences on an outpatient basis can reduce the therapeutic burden and can be a treatment option (85) (LE: 3).

### 5.11 Bladder- and prostatic urethral biopsies

CIS can present as a velvet-like, reddish area that is indistinguishable from inflammation, or it might not be visible at all.

When abnormal areas of urothelium are seen, it is advised to take cold-cup biopsies or biopsies with a resection loop. Biopsies from normal-looking mucosa, so-called random (mapping) biopsies, should be performed in patients with positive urinary cytology and absence of visible bladder tumour, in addition to upper tract work-up/diagnostics. It is recommended to take biopsies from the trigone, bladder dome, and from the right, left, anterior and posterior bladder walls.

In patients with TaT1 tumours, mapping/random biopsies are not routinely recommended. The likelihood of detecting CIS, especially in low-risk tumours, is extremely low (< 2%) (86) (LE: 2a). Material obtained by random or directed biopsies must be sent for pathological assessment in separate containers as specified previously.

Involvement of the prostatic urethra and ducts in men with NMIBC has been reported. Palou *et al.* (87) showed that in 128 men with T1G3 BC, the incidence of CIS in prostatic urethra was 11.7% (LE:2b). The risk of prostatic urethra or duct involvement seems to be higher if the tumour is located on the trigone or bladder neck, in the presence of bladder CIS and multiple tumours (88,89) (LE: 3). When bladder CIS is suspected, or cytology is positive with no evidence of bladder tumour, or abnormalities of prostatic urethra are visible, prostatic urethral biopsies are recommended (87). The biopsy is taken from abnormal areas and from the precollicular area (between 5 and 7 o'clock positions) using a resection loop. In primary NMIBC when stromal invasion is not suspected, a cold-cup biopsy with forceps can be performed (90).

### 5.12 New TURB techniques

#### 5.12.1 New resection techniques

Compared to monopolar resection, the bipolar electrocautery system may reduce the risk of complications (e.g., bladder perforation due to obturator nerve stimulation) (91) (LE: 3). This benefit however must be confirmed by a prospective trial.

#### 5.12.2 New methods of tumour visualisation

As a standard procedure, cystoscopy and TUR are performed using white light. However, the use of white light can lead to missing lesions that are present but not visible, which is why new technologies are being developed.

##### 5.12.2.1 Photodynamic diagnosis (fluorescence cystoscopy)

Photodynamic diagnosis (PDD) is performed using violet light after intravesical instillation of 5-aminolaevulinic acid (ALA) or hexaminolaevulinic acid (HAL). It has been confirmed that fluorescence-guided biopsy and resection are more sensitive than conventional procedures for detection of malignant tumours, particularly for CIS (92,93) (LE: 2a). In the systematic review and meta-analysis, PDD had higher sensitivity than white light endoscopy in the pooled estimates for both patient (92% versus 71%) and biopsy (93% versus 65%) level analyses (93).

PDD had lower specificity than white light endoscopy (63% versus 81%) (93). False-positivity can be induced by inflammation or recent TUR, and during the first 3 months after BCG instillation (94,95) (LE: 3).

Prospective randomised studies evaluating the impact of ALA fluorescence-guided TURB on disease recurrence rate have shown controversial results (93,96,97).

A large, multicentre, prospective randomised trial that compared HAL fluorescence-guided TURB with standard TURB reported an absolute reduction of no more than 9% in the recurrence rate within 9 months in the HAL arm. Median time to recurrence improved from 9.4 months in the white light arm to 16.4 months in the HAL arm after mean follow-up of 53 and 55 months, respectively (98,99) (LE: 1b).

The value of fluorescence cystoscopy for improvement of the outcome in relation to progression rate or survival remains to be demonstrated.

In summary, PDD improves tumour detection rate, particularly in CIS. HAL but not ALA fluorescence-guided TURB was shown to have a beneficial effect on disease recurrence rate.

Photodynamic diagnosis is recommended in patients who are suspected of harbouring a high-grade tumour, for example, for biopsy guidance in patients with positive cytology or with a history of high-grade tumour. The additional costs of the equipment and instillation for PDD should be taken into account.

#### 5.12.2.2 *Narrow band imaging*

In narrow band imaging (NBI) the contrast between normal urothelium and hypervascular cancer tissue is enhanced by filtering white light into two bandwidths of 415 and 540 nm, which are absorbed by haemoglobin. Initial studies have demonstrated improved cancer detection by NBI-guided biopsies and resection (100) (LE: 3). These findings should be confirmed in large multi-institutional studies.

### 5.13 **Second resection**

The significant risk of residual tumour after initial TURB of Ta, T1 lesions has been demonstrated (82,101) (LE: 2a). Persistent disease after resection of T1 tumours has been observed in 33-53% of patients (101-106).

Moreover, the tumour is often understaged by initial resection. The likelihood that a T1 tumour has been understaged and muscle-invasive disease detected by second resection ranges from 4 to 25%. This risk has increased up to 50% in some radical cystectomy series, although these studies have only enrolled selected patients (102,107-109) (LE: 2a). Treatment of a TaT1 high-grade tumour and a T2 tumour is completely different; therefore, correct staging is important.

It has been demonstrated that a second TURB can increase the recurrence-free survival (104,105) (LE: 2a).

A second TURB is recommended in the following situations:

- After incomplete initial TUR;
- If there was no muscle in the specimen after initial resection, with exception of Ta G1 tumours and primary CIS;
- In all T1 tumours;
- In all G3 tumours, except primary CIS.

There is no consensus about the strategy and timing of second TURB. Most authors recommend resection at 2-6 weeks after initial TURB. The procedure should include resection of the primary tumour site.

### 5.14 **Pathological report**

Pathological investigation of the specimen obtained by TURB and biopsies is an essential step in the diagnosis and treatment decision making of bladder cancer.

High quality of resected and submitted tissue is essential for correct pathological assessment. The presence of sufficient muscle is necessary for correct assignment of pT category. Individual biopsies and portions of the tumour should be submitted in separate containers, and labelled individually. Pathologists should obtain from urologists order forms with sufficient clinical information regarding each sample, including the location of each sample.

The pathological report should specify (110):

- location of the evaluated sample (information obtained from the urologist order form);
- grade of each lesion;
- depth of tumour invasion (stage);
- presence of CIS;
- presence of detrusor muscle in the specimen;
- presence of lymphovascular invasion (LVI);
- presence of aberrant histology.

Close cooperation between urologists and pathologists is recommended.

## 5.15 Guidelines for primary assessment of non-muscle-invasive bladder cancers

<b>Initial diagnosis</b>	<b>GR</b>
Patient history should be taken and recorded regarding all important information with possible connection to bladder cancer, including risk factors and history of suspect symptoms.	A
Renal and bladder US may be used during initial work-up in patients with haematuria.	C
At the time of initial diagnosis of bladder cancer, CT urography or IVU should be performed only in selected cases (e.g., tumours located in the trigone).	B
Cystoscopy is recommended in all patients with symptoms suggestive of bladder cancer. It cannot be replaced by cytology or any other non-invasive test.	A
Cystoscopy should describe all macroscopic features of the tumour (site, size, number and appearance) and mucosal abnormalities. A bladder diagram is recommended.	C
Voided urine cytology is advocated to predict high-grade tumour before TUR.	C
Cytology should be performed on fresh urine with adequate fixation. Morning urine is not suitable because of the frequent presence of cytolysis.	C
<b>TURB</b>	
TURB should be performed systematically in individual steps: bimanual palpation under anaesthesia; insertion of the resectoscope, under visual control with inspection of the whole urethra; inspection of the whole urothelial lining of the bladder; biopsy from prostatic urethra (if indicated); cold-cup bladder biopsies (if indicated); resection of the tumour; bimanual palpation after resection; protocol formulation; formulation of order form for pathological evaluation.	C
Perform resection in one piece for small papillary tumours (< 1 cm), including part from the underlying bladder wall.	B
Perform resection in fractions (including muscle tissue) for tumours > 1 cm in diameter.	B
Biopsies should be taken from abnormal-looking urothelium. Biopsies from normal-looking mucosa (trigone, bladder dome, and right, left, anterior and posterior bladder walls) are recommended only when cytology is positive or when exophytic tumour has a non-papillary appearance.	C
Biopsy of the prostatic urethra is recommended for cases of bladder neck tumour, when bladder CIS is present or suspected, when there is positive cytology without evidence of tumour in the bladder, or when abnormalities of the prostatic urethra are visible. If biopsy is not performed during the initial procedure, it should be completed at the time of the second resection.	C
Biopsy of the prostatic urethra should be taken from abnormal areas and from the precollicular area (between 5 and 7 o'clock position) using a resection loop. In primary non-muscle-invasive tumours when stromal invasion is not suspected, the cold-cup biopsy with forceps can be used.	C
If equipment is available, fluorescence-guided (PDD) biopsy should be performed instead of random biopsies when bladder CIS or high-grade tumour is suspected (e.g., positive cytology, recurrent tumour with previous history of a high-grade lesion).	B
The specimens from different biopsies and resection fractions must be referred to the pathologist in separate containers and labelled separately.	C
TURB protocol must describe all steps of the procedure, as well as extent and completeness of resection.	C
A second TURB is recommended in the following situations:	A
- after incomplete initial TURB; - if there is no muscle in the specimen after initial resection, with exception of Ta G1 tumours and primary CIS; - in all T1 tumours; - in all G3 tumours, except primary CIS.	
A second TURB should be performed 2-6 weeks after initial resection.	C
<b>Classification and pathological report</b>	
Depth of tumour invasion is classified according to TNM system.	A
For histological classification, 1973 and 2004 WHO grading systems are used. Until the prognostic role of WHO 2004 is validated by more prospective trials, both classifications should be used.	A
Whenever the terminology NMIBC is used in individual cases, the tumour stage and grade should be mentioned.	A
The pathological report should specify tumour location, tumour grade, depth of tumour invasion, presence of CIS, and whether the detrusor muscle is present in the specimen.	A
The pathological report should specify the presence of LVI or aberrant histology.	C



## 6. PREDICTING DISEASE RECURRENCE AND PROGRESSION

### 6.1 Ta,T1 tumours

The classic way to categorise patients with Ta, T1 tumours with or without concomitant CIS is to divide them into risk groups based on prognostic factors derived from multivariate analyses. Using such a technique, it has been proposed to divide patients into low-, intermediate- and high-risk groups (111). When using these risk groups, however, no distinction is usually drawn between the risk of disease recurrence and disease progression. Although prognostic factors may indicate a high risk of recurrence, the risk of progression might still be low, while other tumours might have a high risk of both recurrence and progression.

In order to predict separately the short-term and long-term risks of disease recurrence and progression in individual patients, the European Organization for Research and Treatment of Cancer (EORTC) Genito-Urinary Cancer Group (GUCCG) has developed a scoring system and risk tables (112). The basis for these tables is the EORTC database, which provides individual patient data for 2,596 patients diagnosed with TaT1 tumours, who were randomised in seven EORTC-GUCCG trials. Patients with CIS alone were not included. Seventy-eight percent of patients received intravesical treatment, mostly chemotherapy. However, they did not undergo a second TUR or receive maintenance BCG. The scoring system is based on the six most significant clinical and pathological factors:

- number of tumours;
- tumour size;
- prior recurrence rate;
- T category;
- presence of concurrent CIS;
- tumour grade.

Table 6 illustrates the weights applied to various factors for calculating the total scores for recurrence and progression. Table 7 shows the total scores stratified, as in the original article (112), into four categories that reflect various probabilities of recurrence and progression at 1 and 5 years.

**Table 6: Weighting used to calculate disease recurrence and progression scores**

Factor	Recurrence	Progression
Number of tumours		
Single	0	0
2-7	3	3
≥ 8	6	3
Tumour diameter		
< 3 cm	0	0
≥ 3 cm	3	3
Prior recurrence rate		
Primary	0	0
≤ 1 recurrence/year	2	2
> 1 recurrence/year	4	2
Category		
Ta	0	0
T1	1	4
Concurrent CIS		
No	0	0
Yes	1	6
Grade (WHO 1973)		
G1	0	0
G2	1	0
G3	2	5
Total score	0-17	0-23

**Table 7: Probability of recurrence and disease progression according to total score**

Recurrence score	Probability of recurrence at 1 year		Probability of recurrence at 5 years	
	%	(95% CI)	%	(95% CI)
0	15	(10-19)	31	(24-37)
1-4	24	(21-26)	46	(42-49)
5-9	38	(35-41)	62	(58-65)
10-17	61	(55-67)	78	(73-84)

Progression score	Probability of progression at 1 year		Probability of progression at 5 years	
	%	(95% CI)	%	(95% CI)
0	0.2	(0-0.7)	0.8	(0-1.7)
2-6	1	(0.4-1.6)	6	(5-8)
7-13	5	(4-7)	17	(14-20)
14-23	17	(10-24)	45	(35-55)

Note: Electronic calculators for Tables 6 and 7, which have been updated for the iPhone, iPad and Android phones and tablets, are available at <http://www.eortc.be/tools/bladdercalculator/>.

A scoring model for BCG-treated patients that predicts the short- and long-term risks of recurrence and progression has been published by the Club Urológico Español de Tratamiento Oncológico (CUETO) (Spanish Urological Oncology Group). It is based on an analysis of 1,062 patients from four CUETO trials that compared different intravesical BCG treatments. Patients received 12 instillations during 5-6 months. No immediate postoperative instillation or second TUR was performed in these patients. The scoring system is based on evaluation of seven prognostic factors:

- sex;
- age;
- prior recurrence status;
- number of tumours;
- T category;
- associated CIS;
- tumour grade.

Using these tables, the calculated risk of recurrence is lower than that obtained by the EORTC tables. For progression, probability is lower only in high-risk patients (113). The lower risks in the CUETO tables may be attributed to using BCG, which is a more effective instillation therapy. Validations of the EORTC scoring system using the CUETO patients treated with BCG and in an independent patient population with long-term follow-up has confirmed its prognostic value (114,115) (LE: 2a).

Further prognostic factors have been described in selected patient populations. Female sex and CIS in the prostatic urethra are important prognostic factors in T1G3 patients treated with an induction course of BCG (87) (LE: 2b). Recurrence at 3 months was the most important predictor of progression in T1G2 tumours treated with TURB (116) (LE: 2b). The prognostic value of pathological factors, particularly T1 substaging, has been discussed elsewhere (see Chapter 4.4).

More work is required to determine the role of molecular markers in improving the predictive accuracy of the currently existing risk tables (114,117).

## 6.2 Carcinoma *in situ*

Without any treatment, approximately 54% of patients with CIS progress to muscle-invasive disease (118) (LE: 3). Unfortunately, there are no reliable prognostic factors that can be used to predict the course of the disease and specify the most dangerous cases. Publications are based on retrospective analyses of small series of patients, and their conclusions are not homogeneous. Some studies have reported a worse prognosis in concurrent CIS and T1 tumours compared to primary CIS (119,120), in extended CIS (121) and in CIS in the prostatic urethra (87) (LE: 3).

Various publications have shown that the response to intravesical treatment with BCG or chemotherapy is an important prognostic factor for subsequent progression and death caused by bladder cancer (113-116). Approximately 10-20% of complete responders eventually progress to muscle-invasive disease, compared with 66% of non-responders (122-124) (LE: 2a).

### 6.3 Recommendation for patients' stratification in risk groups

Based on available prognostic factors and particularly data from the EORTC risk tables, the Guidelines Panel recommends stratification of patients into three risk groups that will facilitate treatment recommendations. Their definition, which takes into account the EORTC risk tables probabilities of recurrence and especially progression, can be found in Table 8. The recommendation is similar but not identical to that provided by the International Bladder Cancer Group (125).

For individual prediction of the risk of tumour recurrence and progression at different intervals after TURB, application of EORTC risk tables and calculator is strongly recommended.

**Table 8: Risk group stratification**

Low-risk tumours	Primary, solitary, Ta, G1 (low grade), < 3 cm, no CIS
Intermediate-risk tumours	All tumours not defined in the two adjacent categories (between the category of low and high risk)
High-risk tumours	Any of the following: <ul style="list-style-type: none"><li>• T1 tumour</li><li>• G3 (high grade) tumour</li><li>• CIS</li><li>• Multiple and recurrent and large (&gt; 3 cm) Ta G1G2 tumours (all conditions must be presented in this point)</li></ul>

#### 6.3.1 Recommendations for stratification of NMIBC

	GR
Stratify patients into three risk groups according to Table 8.	B
Application of EORTC risk tables and calculator for individual prediction of the risk of tumour recurrence and progression in different intervals after TURB.	B

## 7. ADJUVANT TREATMENT

### 7.1 Intravesical chemotherapy

Although state-of-the-art TUR by itself can eradicate a Ta, T1 tumour completely, these tumours commonly recur and can progress to MIBC. The high variability in the 3-month recurrence rate indicates that TUR is incomplete or provokes recurrences in a high percentage of patients (82). It is therefore necessary to consider adjuvant therapy in all patients.

#### 7.1.1 One, immediate, postoperative intravesical instillation of chemotherapy

Early single instillation has been shown to function by the destruction of circulating tumour cells resulting from TUR, and by an ablative effect (chemoresection) on residual tumour cells at the resection site and on small overlooked tumours (126-129) (LE: 3).

In a meta-analysis of 1,476 patients, one immediate instillation of chemotherapy after TUR significantly reduced recurrence rate by 11.7% compared to TUR alone (130) (LE: 1a). The majority of patients (> 80%) in the meta-analysis had a single tumour. A similar efficacy was reported in two more recent studies (131,132), with subgroup analyses suggesting that immediate instillation is most effective in tumour types with the lowest tendency towards recurrence, that is, in single primary or small tumours. Mitomycin C, epirubicin, and doxorubicin have all shown a beneficial effect, with no efficacy comparisons made between the drugs (130) (LE: 1a).

There is evidence from one subgroup analysis and one combined analysis that immediate instillation might have an impact on recurrence even when further adjuvant instillations are given (133,134) (LE: 2a). In contrast, a sufficient number of delayed repeat chemotherapy instillations can also reduce recurrence stemming from tumour implantation (128,133,134). Nevertheless, it is likely that immediate instillation is more effective in preventing recurrence than any of the individual instillations that follow the immediate instillation (128,135) (LE: 3). Clearly, more studies comparing immediate and delayed start regimens are needed.

Prevention of tumour cell implantation should be initiated within the first hours after cell seeding. Within a few hours, the cells are implanted firmly and are covered by extracellular matrix (127,136-138) (LE:

3). In all single instillation studies, the instillation was administered within 24 h. To maximise the efficacy of immediate instillation, one should devise flexible practices that allow the instillation to be given as early as possible, that is, in the recovery room or even in the operating theatre.

Immediate instillation of postoperative chemotherapy should be omitted in any case of overt or suspected intra- or extraperitoneal perforation, which is most likely to appear in extensive TUR procedures, and in situations with bleeding that requires bladder irrigation. Clear instructions should be given to the nursing staff to control the free flow of the bladder catheter at the end of the instillation. Severe complications have been reported in patients with drug extravasation (139).

### 7.1.2 **Additional adjuvant intravesical chemotherapy instillations**

The need for further adjuvant intravesical therapy depends on prognosis. In low-risk patients (Tables 7 and 8), a single immediate instillation reduces the risk of recurrence and is considered as the standard treatment (130) (LE: 1a). No further treatment should be given in these patients before subsequent recurrence. For other patients, however, a single immediate instillation remains an incomplete treatment because of the considerable likelihood of recurrence and/or progression (Tables 7 and 8).

A large meta-analysis of 3,703 patients from 11 randomised trials showed a highly significant 44% reduction in the odds of recurrence at 1 year in favour of chemotherapy over TUR alone (140). This corresponds to an absolute difference of 13-14% in the number of patients with recurrence. Contrary to chemotherapy, two meta-analyses have demonstrated that BCG therapy may reduce the risk tumour progression (141,142) (LE: 1a) (see section 7.2.1). Moreover, BCG maintenance therapy appears to be significantly better in preventing recurrences than regimens with mitomycin C (MMC) or epirubicin (143-145) (see section 7.2.1) (LE: 1a). However, BCG causes significantly more side effects than chemotherapy does (145) (LE: 1a).

It is still controversial for how long and how frequently chemotherapy instillations should be given. From a systematic review of the literature of RCTs, which compared different schedules of intravesical chemotherapy instillations, one can only conclude that the ideal duration and intensity of the schedule remains undefined because of conflicting data (146). The available evidence does not support any treatment longer than 1 year (LE: 3).

### 7.1.3 **Options for improving efficacy of intravesical chemotherapy**

Some promising data have been presented about enhancing the efficacy of MMC using microwave-induced hyperthermia (Synergo) or electromotive drug administration (EMDA) in patients with high-risk tumours. The current evidence, however, is limited. The number of patients in the prospective series applying the microwave-induced hyperthermia was small and data on progression were inconclusive. In one study of 212 patients comparing BCG with sequential BCG and electromotive MMC, a significant benefit was found in favour of the electromotive arm regarding recurrence and progression (147,148) (LE: 2b). Still, both treatment modalities are considered to be experimental.

One RCT using MMC has demonstrated that adapting urinary pH, decreasing urinary excretion, and buffering the intravesical solution reduces the recurrence rate (149) (LE: 1b). Another trial reported that a 1-h instillation of MMC is better than 30 min, but no efficacy comparisons are available for 1- and 2-h instillations (150) (LE: 3).

Another RCT using epirubicin has documented that concentration is more important than treatment duration (151) (LE: 1b). In view of these data, which need confirmation, it seems advisable to ask the patient not to drink on the morning before instillation, and to dissolve the drug in a buffered solution at optimal pH.

## 7.2 **Intravesical Bacillus Calmette-Guérin (BCG) immunotherapy**

### 7.2.1 **Efficacy of BCG**

Five meta-analyses have confirmed that BCG after TUR is superior to TUR alone or TUR and chemotherapy for prevention of recurrence of non-muscle-invasive tumours (143,152-155) (LE: 1a). Three recent RCTs of intermediate- and high-risk tumours have been conducted. BCG was compared with combination of epirubicin and interferon (156), MMC (157) or epirubicin alone (144). All of these studies have confirmed the superiority of BCG for prevention of tumour recurrence (LE: 1a). It has been shown that the effect was long lasting (144,157) and was also observed in a separate analysis of patients with intermediate-risk tumours (144).

One meta-analysis (143) has evaluated the individual data from 2,820 patients enrolled in nine RCTs that have compared MMC versus BCG. In the trials with BCG maintenance, a 32% reduction in the risk of recurrence for BCG compared to MMC was found ( $P < 0.0001$ ), whereas there was a 28% increase in the risk of recurrence ( $P = 0.006$ ) for patients treated with BCG in the trials without BCG maintenance.

Two meta-analyses have demonstrated that BCG therapy prevents, or at least delays, the risk of tumour progression (141,142) (LE: 1a). A meta-analysis carried out by the EORTC-GUCG has evaluated data from 4,863 patients (81.6% with papillary tumours and 18.4% with primary or concurrent CIS) enrolled in 24

RCTs. Five different BCG strains were used, and in 20 of the trials, some form of BCG maintenance was used. Based on a median follow-up of 2.5 years, in 260 out of 2,658 patients (9.8%) treated with BCG, tumours progressed compared to 304 out of 2,205 (13.8%) in the control groups (TUR alone, TUR plus intravesical chemotherapy, or TUR plus other immunotherapy). This shows a reduction of 27% in the odds of progression with BCG maintenance treatment ( $P = 0.0001$ ). The size of the reduction was similar in patients with TaT1 papillary tumours and in those with CIS (142). A recent RCT with long-term observation has demonstrated significantly fewer distant metastases and better overall- and disease-specific survival in patients treated with BCG compared to epirubicin (144) (LE: 1b). On the contrary, a meta-analysis of individual patient data was not able to confirm any statistically significant difference between MMC and BCG for progression, survival and cause of death (143).

The conflicting results in the outcomes of the studies can be explained by different patient characteristics, duration of follow-up, methodology and statistical power. The majority of studies were however able to show a reduction in the risk of progression in high- and intermediate-risk tumours if BCG was applied including a maintenance schedule.

Two other meta-analyses have suggested a possible bias in favour of BCG by the inclusion of patients who were previously treated with intravesical chemotherapy (158,159). In the most recent meta-analysis, however, BCG maintenance was more effective than MMC also in patients who were previously treated with chemotherapy (143) (LE: 1a).

### 7.2.2 **Optimal BCG schedule**

Induction BCG instillations are classically given according to the empirical 6-weekly schedule that was introduced by Morales in 1976 (160). For optimal efficacy, BCG must be given in a maintenance schedule (141-143,155) (LE: 1a). In the EORTC-GU group meta-analysis, only patients who received maintenance BCG benefited. Many different maintenance schedules have been used, ranging from a total of 10 instillations given in 18 weeks, to 27 over 3 years (161). The meta-analysis was unable to determine which BCG maintenance schedule was the most effective (142). In their meta-analysis, Böhle et al. concluded that at least 1 year of maintenance BCG is required to obtain superiority of BCG over MMC for prevention of recurrence or progression (141,155) (LE: 1a).

The optimal number of induction instillations and optimal frequency and duration of maintenance instillations remain unknown (162). However, in an RCT of 1,355 patients, the EORTC has recently shown that when BCG is given at full dose, 3 years maintenance reduces the recurrence rate as compared to 1 year in high-risk but not in intermediate-risk patients. There were no differences in progression or overall survival (159) (LE: 1b). The benefit of the two additional years of maintenance in the high-risk patients should be weighed against its added costs and inconveniences.

### 7.2.3 **BCG toxicity**

BCG intravesical treatment is associated with more side effects compared to intravesical chemotherapy (145) (LE: 1a). Serious side effects however are encountered in < 5% of patients and can be treated effectively in almost all cases (163) (LE: 1b). It has been shown that maintenance schedule is not associated with increased risk of side effects comparing to induction course (163).

Major complications can appear after systemic absorption of the drug. Thus, contraindications of BCG intravesical instillation should be respected.

BCG should not be administered (absolute contraindications):

- during the first 2 weeks after TUR;
- in patients with macroscopic haematuria;
- after traumatic catheterisation;
- in patients with symptomatic urinary tract infection.

The presence of leukocyturia or asymptomatic bacteriuria is not a contraindication for BCG application, and antibiotic prophylaxis is not necessary in these cases (164,165) (LE: 3).

BCG should be used with caution (relative contraindication) in immunocompromised patients (immunosuppression, human immunodeficiency virus (HIV) infection) (166), although some small studies have shown similar efficacy and no increase in complications comparing to non-immuno-compromised patients (167,168) (LE: 3).

The management of side effects after BCG should reflect their type and grade. Recommendations for individual situations have been provided by the International Bladder Cancer Group (IBCG) and by a Spanish group (169,170) (Table 9).

**Table 9: Management options for side effects associated with intravesical BCG (169,171,172)**

<b>Management options for local side effects (modified from IBCG group)</b>	
Symptoms of cystitis	Phenazopyridine, propantheline bromide, or NSAIDs
	If symptoms improve within a few days: continue instillations
	If symptoms persist or worsen: <ol style="list-style-type: none"> <li>Postpone the instillations</li> <li>Perform a urine culture</li> <li>Start empirical antibiotic treatment</li> </ol>
	If symptoms persist even with antibiotic treatment: <ol style="list-style-type: none"> <li>With positive culture: antibiotic treatment according to sensitivity.</li> <li>With negative culture: quinolones and potentially analgesic anti-inflammatory instillations once daily for 5 days (to repeat cycle if necessary) (173).</li> </ol>
	If symptoms persist: anti-tuberculosis drugs + corticosteroids.
	If no response to treatment and/or contracted bladder: radical cystectomy.
Haematuria	Perform urine culture to exclude haemorrhagic cystitis, if other symptoms present.
	If haematuria persists, perform cystoscopy to evaluate presence of bladder tumour.
Symptomatic granulomatous prostatitis	Symptoms rarely present: perform urine culture.
	Quinolones.
	If quinolones are not effective: isoniazid (300 mg/day) and rifampicin (600 mg/day) for 3 months.
	Cessation of intravesical therapy.
Epididymo-orchitis (172)	Perform urine culture and administer quinolones.
	Cessation of intravesical therapy.
	Orchidectomy if abscess or no response to treatment.
<b>Management options for systemic side effects</b>	
General malaise, fever	Generally resolve within 48 h, with or without antipyretics.
Arthralgia and/or arthritis	Rare complication and considered autoimmune reaction.
	Arthralgia: treatment with NSAIDs.
	Arthritis: NSAIDs and if no/partial response proceed to corticosteroids, high-dose quinolones or antituberculous drugs (171).
Persistent high-grade fever (> 38.5°C for > 48 h)	Permanent discontinuation of BCG instillations.
	Immediate evaluation: urine culture, blood tests, chest X-ray.
	Prompt treatment with ≥ 2 antimicrobial agents while diagnostic evaluation is conducted.
	Consultation with an infectious diseases specialist.
BCG sepsis	Prevention: initiate BCG at least 2 weeks post TURBT (if no signs and symptoms of haematuria).
	Cessation of BCG
	For severe infection: <ul style="list-style-type: none"> <li>- High-dose quinolones or isoniazid, rifampicin and ethambutol 1.2 g daily for 6 months.</li> <li>- Early, high-dose corticosteroids as long as symptoms persist.</li> </ul> Consider an empirical non-specific antibiotic to cover Gram-negative bacteria and/or Enterococcus.
Allergic reactions	Antihistamines and anti-inflammatory agents.
	Consider high-dose quinolones or isoniazid and rifampicin for persistent symptoms.
	Delay therapy until reactions resolve.

#### 7.2.4 **Optimal dose of BCG**

To reduce BCG toxicity, instillation of a reduced dose was proposed.

Comparing one-third dose to full-dose BCG in 500 patients, CUETO found no overall difference in efficacy. However, it has been suggested that a full dose of BCG is more effective in multifocal tumours (174, 175) (LE: 1b). Although fewer patients have reported toxicity with the reduced dose, the incidence of severe systemic toxicity was similar in the standard- and reduced-dose groups. The same Spanish group has shown in a prospective RCT that one-third of the standard dose of BCG might be the minimum effective dose for intermediate-risk tumours. A further reduction to one-sixth dose resulted in a decrease in efficacy for prevention of recurrence with no decrease in toxicity (176) (LE: 1b).

The EORTC did not find any difference in toxicity between one-third and full dose BCG, however, the former was associated with a higher recurrence rate, especially when it was given only for 1 year (159) (LE: 1b).

#### 7.2.5 **BCG strain**

There is no conclusive evidence that there may be a difference in clinical efficacy between various BCG strains.

#### 7.2.6 **Indications for BCG**

Although BCG is a very effective treatment, there is a consensus that not all patients with NMIBC should be treated with BCG due to the risk of toxicity. Ultimately, the choice of treatment depends upon the patient's risk (Table 8):

- BCG does not alter the natural course of low-risk tumours (Table 8), and could be considered as overtreatment for this patient category.
- In patients with high-risk tumours, for whom radical cystectomy is not carried out, 1-3 years full-dose maintenance BCG is indicated. The additional beneficial effect of the second and third years of maintenance on recurrence in high-risk patients should be weighed against its added costs and inconveniences.
- In intermediate-risk patients, full-dose BCG with 1 year maintenance is more effective than chemotherapy for prevention of recurrence; however, it has more side effects than chemotherapy. For this reason both BCG with maintenance and intravesical chemotherapy remain an option. The final choice should reflect the individual patient's risk of recurrence and progression as well as efficacy and side effects of each treatment modality.

### 7.3 **Specific aspects of treatment of CIS**

#### 7.3.1 **Treatment strategy**

If concurrent CIS is found in association with MIBC, therapy is determined according to the invasive tumour. The detection of CIS with TaT1 tumours increases the risk of recurrence and progression of TaT1 tumours (112,113) and further treatment is mandatory. The treatment strategy is generally based on the criteria that are summarised in Chapters 7.1, 7.2, 7.4 and 8.

CIS cannot be cured by an endoscopic procedure alone. Histological diagnosis of CIS must be followed by further treatment, either intravesical BCG instillations or radical cystectomy (LE: 4). No consensus exists about whether conservative therapy (intravesical BCG instillations) or aggressive therapy (radical cystectomy) should be done. There has been a lack of randomised trials of instillation therapy and early radical cystectomy as immediate primary treatment. Tumour-specific survival rates after early radical cystectomy for CIS are excellent, but as many as 40-50% of patients might be overtreated (177) (LE: 3).

#### 7.3.2 **Cohort studies**

In retrospective evaluations of patients with CIS, a complete response rate of 48% was achieved with intravesical chemotherapy and 72-93% with BCG (118-121,178) (LE: 2a). Up to 50% of complete responders might eventually show recurrence with a risk of invasion and/or extravesical recurrence (121,161,178,179) (LE: 3).

#### 7.3.3 **Prospective randomised trials**

Unfortunately, there have been few randomised trials in patients with CIS alone. Thus, the power to detect differences in treatment results has been low and the reliability of the conclusions is limited (177).

A meta-analysis of clinical trials that has compared intravesical BCG to intravesical chemotherapy (MMC, epirubicin, or adriamycin) in patients with CIS has shown a significantly increased response rate after BCG and a reduction of 59% in the odds of treatment failure with BCG (OR = 0.41,  $P = 0.0001$ ). In trials that have compared BCG with MMC, the long-term benefit of BCG was smaller, but BCG was superior to MMC in trials with BCG maintenance (OR = 0.57,  $P = 0.04$ ) (180) (LE: 1a).

In an EORTC-GUCG meta-analysis of tumour progression, in a subgroup of 403 patients with CIS, BCG reduced the risk of progression by 35% as compared to intravesical chemotherapy or different

immunotherapy (OR = 0.65, 95% CI = 0.36-1.16,  $P = 0.10$ ) (142) (LE: 1b). There has been no single trial that has demonstrated superiority of combined BCG and MMC over BCG alone (181).

In summary, compared to chemotherapy, BCG treatment of CIS increases the complete response rate, the overall percentage of patients who remain disease free, and reduces the risk of tumour progression (LE: 1a).

#### 7.3.4 Treatment of extravesical CIS

Patients with CIS are at high risk of extravesical involvement: in the upper urinary tract and in the prostatic urethra. Solsona et al. have found that 87 of 138 patients (63%) with CIS developed extravesical involvement initially or during follow-up (182). Patients with extravesical involvement had worse survival than those with bladder CIS alone (182) (LE: 3).

In the prostate, CIS might be present only in the epithelial lining of the prostatic urethra or in the prostatic ducts (177). These situations should be distinguished from tumour invasion into the prostatic stroma, which is staged as T4a, and for which immediate radical cystoprostatectomy is mandatory.

Patients with CIS in the epithelial lining of the prostatic urethra can be treated by intravesical instillation of BCG. TUR of the prostate can improve contact of BCG with the prostatic urethra (78,177,183) (LE: 3).

In patients with prostatic duct involvement, there are promising results, but only from short series, so the data are insufficient to provide clear treatment recommendations (184). No conclusive results have been obtained with conservative therapy, therefore, radical surgery should be considered (183) (LE: 3).

Treatment of CIS that involves the upper urinary tract is discussed in the Guidelines on Urothelial Carcinomas of the Upper Urinary Tract.

## 7.4 Treatment of failure of intravesical therapy

### 7.4.1 Failure of intravesical chemotherapy

Patients with non-muscle-invasive recurrence of BC after chemotherapy regimen can profit from BCG instillations. Prior intravesical chemotherapy has no impact on the effect of BCG instillation (143) (LE: 1a).

### 7.4.2 Recurrence and failure after intravesical BCG immunotherapy

**Table 10: Categories of unsuccessful treatment with intravesical BCG**

<b>BCG failure</b>	
Whenever a muscle-invasive tumour is detected during follow-up.	
BCG-refractory tumour:	
	1. If high-grade, non-muscle-invasive papillary tumour is present at 3 months (185). Further conservative treatment with BCG is connected with increased risk of progression (122,186) (LE: 3).
	2. If CIS (without concomitant papillary tumour) is present at both 3 and 6 months. In patients with CIS present at 3 months, an additional BCG course can achieve a complete response in > 50% of cases (42) LE: 3).
	3. If high-grade tumour appears during BCG therapy.*
High grade recurrence after BCG. Recurrence of high grade/grade 3 (WHO 2004/1973) tumour after completion of BCG maintenance, despite an initial response (187) (LE: 3).*	
<b>BCG intolerance</b>	
Severe side effects that prevent further BCG instillation before completing induction (170).	

\* Patients with low-grade recurrence during or after BCG treatment are not considered as BCG failure.

### 7.4.3 Treatment of BCG failure and recurrences after BCG

Treatment recommendations are provided in Table 11. They reflect categories mentioned in the previous paragraph and tumour characteristics at the time of recurrence.

Patients with BCG failure are unlikely to respond to further BCG therapy; therefore, radical cystectomy is the preferred option. The results of various studies suggests that repeat BCG therapy is appropriate for non-high-grade and even for some high-grade recurrent tumours (188,189) LE: 3).

Additionally, there are now several bladder preservation strategies available that can be categorised as immunotherapy, chemotherapy, device-assisted therapy, and combination therapy (190). Changing from BCG to these options can yield responses in selected cases with BCG treatment failure for NMIBC (188, 191-199) (LE: 3). However, experience is limited and treatments other than radical cystectomy must be considered oncologically inferior at the present time (122,185,186) (LE: 3).



## 7.5 Recommendations for adjuvant therapy in Ta, T1 tumours and for therapy of CIS

	GR
The type of intravesical therapy should be based on the risk groups shown in Tables 8 and 11.	A
One immediate chemotherapy instillation is recommended in tumours presumed to be at low or intermediate risk.	
In patients with low-risk Ta tumours, one immediate instillation of chemotherapy is recommended as the complete adjuvant treatment.	A
In patients with intermediate-risk Ta T1 tumours, one immediate instillation of chemotherapy should be followed by 1 year full-dose BCG treatment, or by further instillation of chemotherapy for a maximum of 1 year.	A
In patients with high-risk tumours, full-dose intravesical BCG for 1-3 years is indicated.	A
In patients with CIS in the epithelial lining of the prostatic urethra, TUR of the prostate followed by intravesical instillation of BCG is an option.	C
In patients at highest risk of tumour progression (Table 11), immediate radical cystectomy should be considered.	C
In BCG refractory tumours, radical cystectomy is indicated.	B
<b>Intravesical chemotherapy.</b>	
One immediate instillation of chemotherapy should be omitted in any case of overt or suspected intra- or extraperitoneal perforation (after extensive TURB, or bleeding requiring bladder irrigation).	C
The optimal schedule of further intravesical chemotherapy instillation and its duration is not defined and should not exceed 1 year.	C
If intravesical chemotherapy is given, it is advised to use the drug at its optimal pH and to maintain the concentration of the drug during instillation by reducing fluid intake.	B
The length of individual instillation should be 1-2 hours.	C
<b>BCG intravesical immunotherapy.</b>	
Absolute contraindications of BCG intravesical instillation are: during the first 2 weeks after TUR; in patients with macroscopic haematuria; after traumatic catheterization; and in patients with symptomatic urinary tract infection.	C
The management of side effects after BCG intravesical instillation should reflect their type and grade (Table 9).	C

## 8. RADICAL CYSTECTOMY FOR NON-MUSCLE-INVASIVE BLADDER CANCER

If cystectomy is indicated before pathologically confirmed progression into muscle-invasive tumour, immediate (immediately after NMIBC diagnosis) and early (after BCG failure) radical cystectomy can be distinguished.

There are several reasons to consider radical cystectomy for selected patients with NMIBC:

- The staging accuracy for T1 tumours by TURB is low with 27-51% of patients being upstaged to muscle-invasive tumour at radical cystectomy (90,108,200-211) (LE: 3).
- Some patients with non-muscle invasive tumours experience disease progression in muscle-invasive disease (Table 7).
- It has been shown retrospectively that patients with high-risk NMIBC who undergo early rather than delayed cystectomy for tumour relapse after initial treatment with TURB and BCG have a better survival rate (212) (LE: 3).

Potential benefit of radical cystectomy must be weighed against the risk, morbidity, and impact on quality of life of radical cystectomy. It is reasonable to propose immediate radical cystectomy to those patients with non-muscle-invasive tumour who are at highest risk of progression. These are (35,87,112,113) (LE: 3):

- multiple and/ or large (> 3 cm) T1, high-grade (G3) tumours;
- T1, high-grade (G3) tumours with concurrent CIS;
- recurrent T1, high-grade (G3) tumours;
- T1G3 and CIS in prostatic urethra;
- micropapillary variant of urothelial carcinoma.

It is recommended to discuss immediate radical cystectomy and conservative treatment with BCG instillation. Patients should be informed about the benefits and risks of both approaches. Individual factors like gender and age of the patient should be considered because of worse prognosis between females and life-long risk of progression after BCG in high risk tumours.

Radical cystectomy is strongly recommended in patients with BCG refractory tumours, as mentioned above. Delay of radical cystectomy might lead to decreased disease-specific survival (213) (LE: 3). In patients in whom radical cystectomy is performed at the time of pathological non-muscle-invasive disease, the 5-year disease-free survival rate exceeds 80% (214-219) (LE: 3).

**Table 11: Treatment recommendations in TaT1 tumours according to risk stratification**

Risk category	Definition	Treatment recommendation
Low-risk tumours	Primary, solitary, Ta, G1, < 3 cm, no CIS	One immediate instillation of chemotherapy
Intermediate-risk tumours	All cases between categories of low and high risk	One immediate instillation of chemotherapy followed by further instillations, either chemotherapy for a maximum of 1 year or 1 year full-dose BCG
High-risk tumours	Any of the following: <ul style="list-style-type: none"> <li>• T1 tumours</li> <li>• G3 tumours</li> <li>• CIS</li> <li>• Multiple <u>and</u> recurrent <u>and</u> large (&gt; 3 cm) Ta G1G2 tumours (all these conditions must be presented)</li> </ul>	Intravesical full-dose BCG instillations for 1-3 years or cystectomy (in highest risk tumours)
Subgroup of highest-risk tumours	T1G3 associated with concurrent bladder CIS, multiple and/or large T1G3 and/or recurrent T1G3, T1G3 with CIS in prostatic urethra, micropapillary variant of urothelial carcinoma	Cystectomy should be considered
	BCG refractory tumours	Cystectomy is recommended

**Table 12: Treatment recommendations for BCG failure and recurrences after BCG**

Category	Treatment recommendation	GR
BCG refractory tumour	1. Radical cystectomy 2. Bladder-preserving strategies in patients not suitable for cystectomy	B
High-grade recurrence after BCG	1. Radical cystectomy 2. Repeat BCG course 3. Bladder-preserving strategies	C
Non-high-grade recurrence after BCG for primary intermediate-risk tumour	1. Repeat BCG or intravesical chemotherapy 2. Radical cystectomy	C

## 9. FOLLOW-UP OF PATIENTS WITH NON-MUSCLE-INVASIVE BLADDER TUMOURS

As a result of the risk of recurrence and progression, patients with TaT1 bladder tumours and with CIS need to be followed up; however, the frequency and duration of cystoscopy and imaging should reflect the individual patient's degree of risk. Using risk tables (see Tables 6 and 7), we are able to predict the short- and long-term risks of recurrence and progression in individual patients, and can adapt the follow-up schedule accordingly (112). When planning the follow-up schedule and methods the following aspects should be considered:

- *The prompt detection of muscle-invasive and high-grade non-muscle-invasive recurrence is crucial because a delay in diagnosis and therapy can be life-threatening.*
- *Tumour recurrence in the low-risk group is nearly always low stage and low grade.*  
Small, non-invasive (Ta), low-grade papillary recurrence does not present an immediate danger to the patient, and early detection is not essential for successful therapy (220-227) (LE: 2b). Fulguration of small papillary recurrences on an outpatient basis could be a safe option that reduces the therapeutic burden (85) (LE: 3). Some authors have even defended temporary surveillance in selected cases (226-228) (LE: 3).
- *The first cystoscopy after TUR at 3 months is a very important prognostic indicator for recurrence and progression (112,116,122,229-231) (LE: 1a).* The first cystoscopy should thus always be performed 3 months after TUR in all patients with TaT1 BC.
- *In tumours at low risk, the risk of recurrence after 5 recurrence-free years is low (230) (LE: 3).* Discontinuation of cystoscopy or its replacement with less-invasive methods can be considered (231).
- *In tumours originally at intermediate or high risk, recurrences after 10 years tumour-free interval are not unusual (232) (LE: 3).* Therefore, lifelong follow-up is recommended (231).
- *The risk of upper urinary tract recurrence increases in patients with multiple and high-risk tumours (47) (LE: 3).*
- *Positive urine test results have a positive impact on the quality of performed follow-up cystoscopy (77) (LE: 1b).* It supports the adjunctive role of urine tests during follow-up.

No non-invasive method has been proposed that can replace endoscopy, therefore, follow-up is based on regular cystoscopy (Section 5.8). There has been a lack of randomised studies that have investigated the possibility of safely reducing the frequency of follow-up cystoscopy. The following recommendations are therefore based mostly on retrospective experience.

### 9.1 Guidelines for follow-up in patients after TURB of NMIBC

	GR
The follow-up of TaT1 tumours is based on regular cystoscopy.	A
Patients with low-risk Ta tumours should undergo cystoscopy at 3 months. If negative, subsequent cystoscopy is advised 9 months later, and then yearly for 5 years.	C
Patients with high-risk tumours should undergo cystoscopy and urinary cytology at 3 months. If negative, subsequent cystoscopy and cytology should be repeated every 3 months for a period of 2 years, and every 6 months thereafter until 5 years, and then yearly.	C
Patients with intermediate-risk TaT1 tumours should have an in-between follow-up scheme using cystoscopy and cytology, which is adapted according to personal and subjective factors.	C
Regular (yearly) upper tract imaging (CT-IVU or IVU) is recommended for high-risk tumours.	C
Endoscopy under anaesthesia and bladder biopsies should be performed when office cystoscopy shows suspicious findings or if urinary cytology is positive.	B
During follow-up in patients with positive cytology and no visible tumour in the bladder, R-biopsies or biopsies with PDD (if equipment is available) and investigation of extravesical locations (CT urography, prostatic urethra biopsy) are recommended.	B

## 10. REFERENCES

1. Ferlay J, S.H., Bray F, Forman D, et al. GLOBOCAN 2008 v1.2, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 10 2010, International Agency for Research on Cancer: Lyon, France.  
<http://www.iarc.fr/en/publications/eresources/cancerbases/index.php>
2. Ploeg M, Aben KKH, Kiemeny LA. The present and future burden of urinary bladder cancer in the world. *World J Urol* 2009 Jun;27(3):289-93.  
<http://www.ncbi.nlm.nih.gov/pubmed/19219610>
3. Burger M, Catto JW, Dalbagni G, et al. Epidemiology and risk factors of urothelial bladder cancer. *Eur Urol* 2013 Feb;63(2):234-41.  
<http://www.ncbi.nlm.nih.gov/pubmed/22877502>
4. Bosetti C, et al. Trends in mortality from urologic cancers in Europe, 1970-2008. *Eur Urol*, 2011. 60(1): p. 1-15.
5. Ferlay J., Randi G, Bosetti C, et al., Declining mortality from bladder cancer in Europe. *BJU Int* 2008 Jan. 101(1):11-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/17971176>
6. Rafnar T, Vermeulen SH, Sulem P, et al. European genome-wide association study identifies SLC14A1 as a new urinary bladder cancer susceptibility gene. *Hum Mol Genet* 2011 Nov 1;20(21):4268-81.  
<http://www.ncbi.nlm.nih.gov/pubmed/21750109>
7. Freedman ND, Silverman DT, Hollenbeck AR, et al. Association between smoking and risk of bladder cancer among men and women. *JAMA* 2011 Aug 17;306(7):737-45.  
<http://www.ncbi.nlm.nih.gov/pubmed/21846855>
8. Rushton L, Bagga S, Bevan R, et al. Occupation and cancer in Britain. *Br J Cancer* 2010 Apr 27;102(9):1428-37.  
<http://www.ncbi.nlm.nih.gov/pubmed/20424618>
9. Rushton L, Huchings SJ, Fortunato L, et al. Occupational cancer burden in Great Britain. *Br J Cancer* 2012 Jun 19;107 Suppl 1:S3-S7.  
<http://www.ncbi.nlm.nih.gov/pubmed/22710676>
10. Samanic CM, Kogevinas M, Silverman DT, et al. Occupation and bladder cancer in a hospital-based case-control study in Spain. *Occup Environ Med* 2008 May;65(5):347-53.  
<http://www.ncbi.nlm.nih.gov/pubmed/17951336>
11. Ros MM, Bas Bueno-de-Mesquita HB, Büchner FL, et al. Fluid intake and the risk of urothelial cell carcinomas in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Int J Cancer* 2011 Jun 1;128(11):2695-708.  
<http://www.ncbi.nlm.nih.gov/pubmed/20715171>
12. Koutros S, Silverman DT, Baris D, et al. Hair dye use and risk of bladder cancer in the New England bladder cancer study. *Int J Cancer* 2011 Dec 15;129(12):2894-904.  
<http://www.ncbi.nlm.nih.gov/pubmed/21678399>
13. Ros MM, Gago-Dominguez M, Aben KK, et al. Personal hair dye use and the risk of bladder cancer: a case-control study from The Netherlands. *Cancer Causes Control* 2012 Jul;23(7):1139-48.  
<http://www.ncbi.nlm.nih.gov/pubmed/22581032>
14. Sobin LH, Gospodariwicz M, Wittekind C (eds). TNM classification of malignant tumors. UICC International Union Against Cancer. 7th edn. Wiley-Blackwell, 2009 Dec; pp. 262-265.  
<http://www.uicc.org/tnm/>
15. Epstein JI, Amin MB, Reuter VR, et al. The World Health Organization/International Society of Urological Pathology consensus classification of urothelial (transitional cell) neoplasms of the urinary bladder. *Am J Surg Pathol* 1998 Dec;22(12):1435-48.  
<http://www.ncbi.nlm.nih.gov/pubmed/9850170>
16. Sauter G, Algaba F, Amin M, et al. Tumours of the urinary system: non-invasive urothelial neoplasias. In: Eble JN, Sauter G, Epstein JI, Sesterhenn I, eds. WHO classification of classification of tumours of the urinary system and male genital organs. Lyon: IARCC Press, 2004, pp. 29-34.
17. Pan CC, Chang YH, Chen KK, et al. Prognostic significance of the 2004 WHO/ISUP classification for prediction of recurrence, progression, and cancer-specific mortality of non-muscle-invasive urothelial tumors of the urinary bladder: a clinicopathologic study of 1,515 cases. *Am J Clin Pathol* 2010 May;133(5):788-95.  
<http://www.ncbi.nlm.nih.gov/pubmed/20395527>
18. Burger M, Catto JW, Dalbagni G, et al. Epidemiology and risk factors of urothelial bladder cancer. *Eur Urol*. 2013 Feb;63(2):234-41.  
<http://www.ncbi.nlm.nih.gov/pubmed/22877502>

19. MacLennan GT, Kirkali Z, Cheng L. Histologic grading of noninvasive papillary urothelial neoplasms. *Eur Urol* 2007 Apr;51(4):889-98.  
<http://www.ncbi.nlm.nih.gov/pubmed/17095142>
20. May M, Brookman-Amisshah S, Roigas J, et al. Prognostic accuracy of individual uropathologists in Non-invasive urinary bladder carcinoma: a multicentre study comparing the 1973 and 2004 World Health Organisation Classifications. *Eur Urol* 2010 May;57(5):850-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/19346063>
21. van Rhijn BW, van Leenders GJ, Ooms BC, et al. The pathologist's mean grade is constant and individualizes the prognostic value of bladder cancer grading. *Eur Urol* 2010 Jun;57(6):1052-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/19765886>
22. Otto W, et al. The WHO classification of 1973 is more suitable than the WHO classification of 2004 for predicting survival in pT1 urothelial bladder cancer. *BJU Int* 2011 Feb;107(3):404-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/20707791>
23. Pellucchi F, Freschi M, Ibrahim B, et al. Clinical reliability of the 2004 WHO histological classification system compared with the 1973 WHO system for Ta primary bladder tumors. *J Urol* 2011 Dec;186(6):2194-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/22019037>
24. Cao D, Vollmer RT, Luly J, et al. Comparison of 2004 and 1973 World Health Organization grading systems and their relationship to pathologic staging for predicting long-term prognosis in patients with urothelial carcinoma. *Urology* 2010 Sep;76(3):593-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/20381131>
25. Murphy WM, Takezawa K, Maruniak NA. Interobserver discrepancy using the 1998 WHO/ISUP classification of urothelial neoplasms: practical choices for patient care. *J Urol* 2002 Sep;168(3):968-72.  
<http://www.ncbi.nlm.nih.gov/pubmed/12187201>
26. Witjes JA, Moonen PMJ, van der Heijden AG. Review pathology in a diagnostic bladder cancer trial. *Urology* 2006 Apr;67(4):751-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/16566990>
27. Bol MG, Baak J, Buhr-Wildhagen S, et al. Reproducibility and prognostic variability of grade and lamina propria invasion in stages Ta, T1 urothelial carcinoma of the bladder. *J Urol* 2003 Apr;169(4):1291-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/12629345>
28. Van der Meijden A, Sylvester R, Collette L, et al. The role and impact of pathology review on stage and grade assessment on stages Ta and T1 bladder tumors: a combined analysis of 5 European Organization for Research and Treatment of Cancer trials. *J Urol* 2000 Nov;164(5):1533-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/11025698>
29. Van Rhijn BWG, van der Kwast TH, Kakiashvili DM, et al. Pathological stage review is indicated in primary pT1 bladder cancer. *BJU Int* 2010 Jul;106(2):206-11.  
<http://www.ncbi.nlm.nih.gov/pubmed/20002439>
30. Orsola A, Trias I, Raventós CX, et al. Initial high-grade T1 urothelial cell carcinoma: feasibility and prognostic significance of lamina propria invasion microstaging (T1a/b/c) in BCG-treated and BCG-non-treated patients. *Eur Urol*. 2005 Aug;48(2):231-8, discussion 238.  
<http://www.ncbi.nlm.nih.gov/pubmed/15963635>
31. Andius P, Johansson SL, Holmäng S. Prognostic factors in stage T1 bladder cancer: tumor pattern (solid or papillary) and vascular invasion more important than depth of invasion. *Urology* 2007 Oct;70(4):758-62.  
<http://www.ncbi.nlm.nih.gov/pubmed/17991551>
32. van Rhijn BW, van der Kwast TH, Alkhateeb SS, et al. A new and highly prognostic system to discern T1 bladder cancer substage. *Eur Urol*. 2012 Feb;61(2):378-84.  
<http://www.ncbi.nlm.nih.gov/pubmed/22036775>
33. van Rhijn BW, Liu L, Vis AN, et al. Prognostic value of molecular markers, sub-stage and European Organisation for the Research and Treatment of Cancer risk scores in primary T1 bladder cancer. *BJU Int* 2012 Oct;110(8):1169-76.  
<http://www.ncbi.nlm.nih.gov/pubmed/22448597>
34. Cho KS, Seo HK, Joung JY et al. Lymphovascular Invasion in Transurethral Resection Specimens as Predictor of Progression and Metastasis in Patients With Newly Diagnosed T1 Bladder Urothelial Cancer. *J Urol* 2009 Dec;182(6):2625-31.  
<http://www.ncbi.nlm.nih.gov/pubmed/19836779>

35. Kamat AM. The Case for Early Cystectomy in the Treatment of Non-muscle Invasive Micropapillary Bladder Carcinoma. *J Urol* 2006 Mar;175(3 Pt 1):881-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/16469571>
36. Amin A, Epstein JI. Noninvasive micropapillary urothelial carcinoma: a clinicopathologic study of 18 cases. *Hum Pathol* 2012 Dec;43(12):2124-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/22939957>
37. Comperat E, Roupret M, Yaxley J, et al. Micropapillary urothelial carcinoma of the urinary bladder: a clinicopathological analysis of 72 cases. *Pathology* 2010 Dec;42(7):650-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/21080874>
38. Blochin EB, Park KJ, Tickoo SK, et al. Urothelial carcinoma with prominent squamous differentiation in the setting of neurogenic bladder: role of human papillomavirus infection. *Mod Pathol* 2012 Nov;25(11):1534-42.  
<http://www.ncbi.nlm.nih.gov/pubmed/22766788>
39. Frstrup N, Ulhøi BP, Birkenkamp-Demtröder K, et al. Cathepsin E, maspin, Plk1, and survivin are promising prognostic protein markers for progression in non-muscle invasive bladder cancer. *Am J Pathol* 2012 May;180(5):1824-34.  
<http://www.ncbi.nlm.nih.gov/pubmed/22449953>
40. Palou J, Algaba F, Vera I, et al. Protein expression patterns of ezrin are predictors of progression in T1G3 bladder tumours treated with nonmaintenance bacillus Calmette-Guérin. *Eur Urol* 2009 Nov;56(5):829-36.  
<http://www.ncbi.nlm.nih.gov/pubmed/18926620>
41. van Rhijn BW, van der Kwast TH, Liu L, et al. The FGFR3 mutation is related to favorable pT1 bladder cancer. *J Urol* 2012 Jan;187(1):310-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/22099989>
42. Sylvester R, van der Meijden A, Witjes JA, et al. High-grade Ta urothelial carcinoma and carcinoma in situ of the bladder. *Urology* 2005 Dec;66(6 Suppl 1):90-107.  
<http://www.ncbi.nlm.nih.gov/pubmed/16399418>
43. Lamm DL, Herr HW, Jakse G, et al. Updated concepts and treatment of carcinoma in situ. *Urol Oncol* Jul-Oct 1998;4(4-5):130-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/21227218>
44. Goessl C, Knispel HH, Millar K, et al. Is routine excretory urography necessary at first diagnosis of bladder cancer? *J Urol* 1997 Feb;157(2):480-1.  
<http://www.ncbi.nlm.nih.gov/pubmed/8996338>
45. Palou J, Rodriguez-Rubio F, Huguet J, et al. Multivariate analysis of clinical parameters of synchronous primary superficial bladder cancer and upper urinary tract tumours. *J Urol* 2005 Sep;174(3):859-61.  
<http://www.ncbi.nlm.nih.gov/pubmed/16093970>
46. Holmang S, Hedelin H, Anderstrom C, et al. Long-term follow-up of a bladder carcinoma cohort: routine followup urography is not necessary. *J Urol* 1998 Jul;160(1):45-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/9628602>
47. Millán-Rodríguez F, Chéchile-Toniolo G, Salvador-Bayarri J, et al. Upper urinary tract tumours after primary superficial bladder tumours: prognostic factors and risk groups. *J Urol* 2000 Oct;164(4):1183-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/10992362>
48. Nolte-Ernsting C, Cowan N. Understanding multislice CT urography techniques: many roads lead to Rome. *Eur Radiol* 2006 Dec;16(12):2670-86.  
<http://www.ncbi.nlm.nih.gov/pubmed/16953373>
49. Têtu B. Diagnosis of urothelial carcinoma from urine. *Mod Pathol* 2009 Jun;22 Suppl 2:S53-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/19494853>
50. Raitanen M-P, Aine R, Rintala E, et al; FinnBladder Group. Differences between local and review urinary cytology and diagnosis of bladder cancer. An interobserver multicenter analysis. *Eur Urol* 2002;41(3):284-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/12180229>
51. Lokeshwar VB, Habuchi T, Grossman HB, et al. Bladder tumour markers beyond cytology: international consensus panel on bladder tumour markers. *Urology* 2005 Dec;66(6 Suppl 1):35-63.  
<http://www.ncbi.nlm.nih.gov/pubmed/16399415>
52. Glas AS, Roos D, Deutekom M, et al. Tumour markers in the diagnosis of primary bladder cancer. A systematic review. *J Urol* 2003 Jun;169(6):1975-82.  
<http://www.ncbi.nlm.nih.gov/pubmed/12771702>

53. Lotan Y, Roehrborn CG. Sensitivity and specificity of commonly available bladder tumour markers versus cytology: results of a comprehensive literature review and meta-analysis. *Urology* 2003 Jan;61(1):109-18.  
<http://www.ncbi.nlm.nih.gov/pubmed/12559279>
54. Van Rhijn BW, van der Poel HG, van der Kwast TH. Urine markers for bladder cancer surveillance: a systematic review. *Eur Urol* 2005 Jun;47(6):736-48.  
<http://www.ncbi.nlm.nih.gov/pubmed/15925067>
55. Vrooman OPJ, Witjes JA. Urinary markers in bladder cancer. *Eur Urol* 2008 May;53(5):909-16.  
<http://www.ncbi.nlm.nih.gov/pubmed/18162285>
56. Lotan Y, Shariat SF, Schmitz-Drager BJ, et al. Considerations on implementing diagnostic markers into clinical decision making in bladder cancer. *Urol Oncol* 2010 Jul-Aug;28(4):441-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/20610281>
57. Van Rhijn BWG, van der Poel HG, van der Kwast HG. Cytology and urinary markers for the diagnosis of bladder cancer. *Eur Urol Suppl* 2009;8:536-41.
58. Catto JWF. Old and new urinary markers: Which one is the PSA for bladder cancer? *Eur Urol Suppl*. 2008;7:422-5.
59. Yutkin V, Nisman B, Pode D. Can urinary biomarkers replace cystoscopic examination in bladder cancer surveillance? *Expert Rev Anticanc* 2010 Jun;10(6):787-90.  
<http://www.ncbi.nlm.nih.gov/pubmed/20553203>
60. Zwarthoff EC. Detection of tumours of the urinary tract in voided urine. *Scand J Urol Nephrol* 2008 Sep;(218):147-53.  
<http://www.ncbi.nlm.nih.gov/pubmed/18815929>
61. Shirodkar SP, Lokeshwar VB. Bladder tumor markers: from hematuria to molecular diagnostics - where do we stand? *Expert Rev Anticanc* 2008 Jul;8(7):1111-23.  
<http://www.ncbi.nlm.nih.gov/pubmed/18588456>
62. Shariat SF, Karam JA, Lerner SP. Molecular markers in bladder cancer. *Curr Opin Urol*. 2008 Jan;18(1):1-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/18090481>
63. Kim WJ, Bae SC. Molecular biomarkers in urothelial bladder cancer. *Cancer Sci* 2008 Apr;99(4):646-52.  
<http://www.ncbi.nlm.nih.gov/pubmed/18377416>
64. Agarwal PK, Black PC, Kamat AM. Considerations on the use of diagnostic markers in management of patients with bladder cancer. *World J Urol* 2008 Feb;26(1):39-44.  
<http://www.ncbi.nlm.nih.gov/pubmed/18092171>
65. Kausch I, Böhle A. Bladder cancer II. Molecular aspects and diagnosis. *Eur Urol* 2001 May;39(5):498-506.  
<http://www.ncbi.nlm.nih.gov/pubmed/11464029>
66. Hajdinjak T. UroVysion FISH test for detecting urothelial cancers: Meta-analysis of diagnostic accuracy and comparison with urinary cytology testing. *Urol Oncol* 2008 Nov-Dec;26(6):645-51.  
<http://www.ncbi.nlm.nih.gov/pubmed/18367109>
67. Schlomer BJ, Ho R, Sagalowsky A, et al. Prospective Validation of the Clinical Usefulness of Reflex Fluorescence In Situ Hybridization Assay in Patients With Atypical Cytology for the Detection of Urothelial Carcinoma of the Bladder. *J Urol* 2010 Jan;183(1):62-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/19913822>
68. Bergman J, Reznicek RC, Rajfer J. Surveillance of patients with bladder carcinoma using fluorescent in-situ hybridization on bladder washings. *BJU Int* 2008 Jan;101(1):26-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/17850364>
69. Van der Aa MN, Zwarthoff EC, Steyerberg EW, et al. Microsatellite Analysis of Voided-Urine Samples for Surveillance of Low-Grade Non-Muscle-Invasive Urothelial Carcinoma: Feasibility and Clinical Utility in a Prospective Multicenter Study (Cost-Effectiveness of Follow-Up of Urinary Bladder Cancer Trial (CEFUB)). *Eur Urol* 2009 Mar;55(3):659-67.  
<http://www.ncbi.nlm.nih.gov/pubmed/18501499>
70. De Bekker-Grob EW, van der Aa MN, Zwarthoff EC, et al. Non-muscle-invasive bladder cancer surveillance for which cystoscopy is partly replaced by microsatellite analysis of urine: a cost-effective alternative? *BJU Int* 2009 Jul;104(1):41-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/19500328>
71. Rouprêt M, Hupertan V, Yates DR, et al. A comparison of the performance of microsatellite and methylation urine analysis for predicting the recurrence of urothelial cell carcinoma, and definition of a set of markers by Bayesian network analysis. *BJU Int* 2008 Jun;101(11):1448-53.  
<http://www.ncbi.nlm.nih.gov/pubmed/18325051>

72. Grossman HB, Messing E, Soloway M, et al, Detection of bladder cancer using a point-of-care proteomic assay, *JAMA* 2005 Feb 293(7):810-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/15713770>
73. Lotan Y, Svatek RS, Malats N. Screening for bladder cancer: a perspective. *World J Urol* 2008 Feb;26(1):13-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/18030473>
74. Roobol MJ, Bangma CH, el Bouazzaoui S, et al. Feasibility study of screening for bladder cancer with urinary molecular markers (the BLU-P project). *Urol Oncol* 2010 Nov-Dec;28(6):686-90.  
<http://www.ncbi.nlm.nih.gov/pubmed/21062653>
75. Grossman HB, Soloway M, Messing E, et al. Surveillance for recurrent bladder cancer using a point-of-care proteomic assay. *JAMA* 2006 Jan;295(3):299-305.  
<http://www.ncbi.nlm.nih.gov/pubmed/16418465>
76. Babjuk M, Soukup V, Pesl M, et al. Urinary cytology and quantitative BTA and UBC tests in surveillance of patients with pT1 bladder urothelial carcinoma. *Urology* 2008 Apr;71(4):718-22.  
<http://www.ncbi.nlm.nih.gov/pubmed/18387400>
77. van der Aa MN, Steyerberg EW, Bangma C, et al. Cystoscopy revisited as the gold standard for detecting bladder cancer recurrence: diagnostic review bias in the randomized, prospective CEFUB trial. *J Urol* 2010 Jan;183(1):76-80.  
<http://www.ncbi.nlm.nih.gov/pubmed/19913254>
78. Kurth KH, Schellhammer PF, Okajima E, et al. Current methods of assessing and treating carcinoma in situ of the bladder with or without involvement of the prostatic urethra. *Int J Urol* 1995 Jun;2(Suppl 2):8-22.  
<http://www.ncbi.nlm.nih.gov/pubmed/7553309>
79. Aaronson DS, Walsh TJ, Smith JF, et al. Meta-analysis: does lidocaine gel before flexible cystoscopy provide pain relief? *BJU Int* 2009 Aug;104(4):506-9; discussion 509-10.  
<http://www.ncbi.nlm.nih.gov/pubmed/19239453>
80. Blick CG, Nazir SA, Mallett S, et al. Evaluation of diagnostic strategies for bladder cancer using computed tomography (CT) urography, flexible cystoscopy and voided urine cytology: results for 778 patients from a hospital haematuria clinic. *BJU Int* 2012 Jul;110(1):84-94.  
<http://www.ncbi.nlm.nih.gov/pubmed/22122739>
81. Richterstetter M, Wullich B, Amann K, et al. The value of extended transurethral resection of bladder tumour (TURBT) in the treatment of bladder cancer. *BJU Int* 2012 Jul;110(2 Pt 2):E76-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/22313727>
82. Brausi M, Collette L, Kurth K, et al; EORTC Genito-Urinary Tract Cancer Collaborative Group. Variability in the recurrence Rate at first follow-up cystoscopy after TUR in stage Ta T1 transitional cell carcinoma of the bladder: a combined analysis of seven EORTC studies. *Eur Urol* 2002 May;41(5): 523-31.  
<http://www.ncbi.nlm.nih.gov/pubmed/12074794>
83. Mariappan P, Zachou A, Grigor KM. Detrusor muscle in the first, apparently complete transurethral resection of bladder tumour specimen is a surrogate marker of resection quality, predicts risk of early recurrence, and is dependent on operator experience. *Eur Urol* 2010 May;57(5):843-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/19524354>
84. Mariappan P, Finney SM, Head E, et al. Good quality white-light transurethral resection of bladder tumours (GQ-WLTURBT) with experienced surgeons performing complete resections and obtaining detrusor muscle reduces early recurrence in new non-muscle-invasive bladder cancer: validation across time and place and recommendation for benchmarking. *BJU Int* 2012 Jun;109(11):1666-73.  
<http://www.ncbi.nlm.nih.gov/pubmed/22044434>
85. Herr HW, Donat SM, Reuter VE. Management of Low Grade Papillary Bladder Tumors. *J Urol* 2007 Oct;178(4 Pt 1):1201-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/17698090>
86. Van der Meijden A, Oosterlinck W, Brausi M, et al. Significance of bladder biopsies in Ta,T1 bladder tumours: a report of the EORTC Genito-Urinary Tract Cancer Cooperative Group. EORTC-GU Group Superficial Bladder Committee. *Eur Urol* 1999 Apr;35(4):267-71.  
<http://www.ncbi.nlm.nih.gov/pubmed/10419345>
87. Palou J, Sylvester RJ, Faba OR, et al. Female gender and carcinoma in situ in the prostatic urethra are prognostic factors for recurrence, progression, and disease-specific mortality in T1G3 bladder cancer patients treated with bacillus Calmette-Guérin. *Eur Urol* 2012 Jul;62(1):118-25.  
<http://www.ncbi.nlm.nih.gov/pubmed/22101115>



88. Matzkin H, Soloway MS, Hardeman S. Transitional cell carcinoma of the prostate. *J Urol* 1991 Nov;146(5):1207-12.  
<http://www.ncbi.nlm.nih.gov/pubmed/1942262>
89. Mungan MU, Canda AE, Tuzel E, et al. Risk factors for mucosal prostatic urethral involvement in superficial transitional cell carcinoma of the bladder. *Eur Urol* 2005 Nov;48(5):760-3.  
<http://www.ncbi.nlm.nih.gov/pubmed/16005563>
90. Huguet J, Crego M, Sabaté S, et al. Cystectomy in patients with high risk superficial bladder tumors who fail intravesical BCG therapy: pre-cystectomy prostate involvement as a prognostic factor. *Eur Urol* 2005 Jul;48(1):53-9; discussion 59.  
<http://www.ncbi.nlm.nih.gov/pubmed/15967252>
91. Gupta NP, Nayyar R, Hemal AK, et al. Outcome analysis of robotic pyeloplasty: a large single-centre experience. *BJU Int* 2010 Apr;105(7):980-3.  
<http://www.ncbi.nlm.nih.gov/pubmed/19874304>
92. Kausch I, Sommerauer M, Montorsi F, et al. Photodynamic diagnosis in non-muscle-invasive bladder cancer: a systematic review and cumulative analysis of prospective studies. *Eur Urol* 2010 Apr;57(4):595-606.  
<http://www.ncbi.nlm.nih.gov/pubmed/20004052>
93. Mowatt G, N'Dow J, Vale L, et al; Aberdeen Technology Assessment Review (TAR) Group. Photodynamic diagnosis of bladder cancer compared with white light cystoscopy: Systematic review and meta-analysis. *Int J Technol Assess Health Care* 2011 Jan;27(1):3-10.  
<http://www.ncbi.nlm.nih.gov/pubmed/21262078>
94. Draga RO, Grimbergen MC, Kok ET, et al. Photodynamic diagnosis (5 aminolevulinic acid) of transitional cell carcinoma after bacillus Calmette- Guérin immunotherapy and mitomycin C intravesical therapy. *Eur Urol* 2010 Apr;57(4):655-60.  
<http://www.ncbi.nlm.nih.gov/pubmed/19819064>
95. Ray ER, Chatterton K, Khan MS, et al. Hexylaminolaevulinate fluorescence cystoscopy in patients previously treated with intravesical bacille Calmette-Guérin. *BJU Int* 2010 Mar;105(6):789-94.  
<http://www.ncbi.nlm.nih.gov/pubmed/19832725>
96. Schumacher MC, Holmäng S, Davidsson T, et al. Transurethral resection of non-muscle-invasive bladder transitional cell cancers with or without 5-aminolevulinic Acid under visible and fluorescent light: results of a prospective, randomised, multicentre study. *Eur Urol* 2010 Feb;57(2):293-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/19913351>
97. Stenzl A, Penkoff H, Dajc-Sommerer E, et al. Detection and clinical outcome of urinary bladder cancer with 5-aminolevulinic acid-induced fluorescence cystoscopy : A multicenter randomized, double-blind, placebo-controlled trial. *Cancer* 2011 Mar 1;117(5):938-47.  
<http://www.ncbi.nlm.nih.gov/pubmed/21351082>
98. Stenzl A, Burger M, Fradet Y, et al. Hexaminolevulinate guided fluorescence cystoscopy reduces recurrence in patients with nonmuscle invasive bladder cancer. *J Urol* 2010 Nov;184(5):1907-13.  
<http://www.ncbi.nlm.nih.gov/pubmed/20850152>
99. Grossman HB, Stenzl A, Fradet Y, et al. Long-term decrease in bladder cancer recurrence with hexaminolevulinate enabled fluorescence cystoscopy. *J Urol* 2012 Jul;188(1):58-62.  
<http://www.ncbi.nlm.nih.gov/pubmed/22583635>
100. Cauberg EC, Kloen S, Visser M, et al. Narrow band imaging cystoscopy improves the detection of non-muscle-invasive bladder cancer. *Urology* 2010 Sep;76(3):658-63  
<http://www.ncbi.nlm.nih.gov/pubmed/20223505>
101. Miladi M, Peyromaure M, Zerbib M, et al. The value of a second transurethral resection in evaluating patients with bladder tumours. *Eur Urol* 2003 Mar;43(3):241-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/12600426>
102. Brauers A, Buettner R, Jakse G. Second resection and prognosis of primary high risk superficial bladder cancer: is cystectomy often too early? *J Urol* 2001 Mar;165(3):808-10.  
<http://www.ncbi.nlm.nih.gov/pubmed/11176474>
103. Schips L, Augustin H, Zigeuner RE, et al. Is repeated transurethral resection justified in patients with newly diagnosed superficial bladder cancer? *Urology* 2002 Feb;59(2):220-3.  
<http://www.ncbi.nlm.nih.gov/pubmed/11834389>
104. Grimm M-O, Steinhoff Ch, Simon X, et al. Effect of routine repeat transurethral resection for superficial bladder cancer: a long-term observational study. *J Urol* 2003 Aug;170(2 Pt 1):433-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/12853793>

105. Divrik RT, Yildirim Ü, Zorlu F, et al. The effect of repeat transurethral resection on recurrence and progression rates in patients with T1 tumours of the bladder who received intravesical mitomycin: a prospective, randomized clinical trial. *J Urol* 2006 May;175(5):1641-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/16600720>
106. Jahnsen S, Wiklund F, Duchek M, et al. Results of second-look resection after primary resection of T1 tumour of the urinary bladder. *Scand J Urol Nephrol* 2005;39(3):206-10.  
<http://www.ncbi.nlm.nih.gov/pubmed/16127800>
107. Dalbagni G, Vora K, Kaag M, et al. Clinical Outcome in a Contemporary Series of Restaged Patients with Clinical T1 Bladder Cancer. *Eur Urol* 2009 Dec;56(6):903-10.  
<http://www.ncbi.nlm.nih.gov/pubmed/19632765>
108. Fritsche HM, Burger M, Svatek RS, et al. Characteristics and outcomes of patients with clinical T1 grade 3 urothelial carcinoma treated with radical cystectomy: results from an international cohort. *Eur Urol* 2010 Feb;57(2):300-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/19766384>
109. Kulkarni GS, Hakenberg OW, Gschwend JE, et al. An updated critical analysis of the treatment strategy for newly diagnosed high-grade T1 (previously T1G3) bladder cancer. *Eur Urol* 2010 Jan;57(1):60-70.  
<http://www.ncbi.nlm.nih.gov/pubmed/19740595>
110. Lopez-Beltran A, Bassi P, Pavone-Macaluso M, et al. Handling and pathology reporting of specimens with carcinoma of the urinary bladder, ureter, and renal pelvis. *Eur Urol* 2004 Mar;45(3):257-66.  
<http://www.ncbi.nlm.nih.gov/pubmed/15036668>
111. Millan-Rodriguez F, Chechile-Toniolo G, Salvador-Bayarri J, et al. Primary superficial bladder cancer risk groups according to progression, mortality and recurrence. *J Urol* 2000 Sep;164(3 Pt 1):680-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/10954628>
112. Sylvester RJ, van der Meijden AP, Oosterlinck W, et al. Predicting recurrence and progression in individual patients with stage TaT1 bladder cancer using EORTC risk tables: a combined analysis of 2596 patients from seven EORTC trials. *Eur Urol* 2006 Mar;49(3):466-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/16442208>
113. Fernandez-Gomez J, Madero R, Solsona E, et al. Predicting nonmuscle invasive bladder cancer recurrence and progression in patients treated with bacillus Calmette-Guerin: the CUETO scoring model. *J Urol* 2009 Nov;182(5):2195-203.  
<http://www.ncbi.nlm.nih.gov/pubmed/19758621>
114. Van Rhijn BW, Zuiverloon TC, Vis AN, et al. Molecular grade (FGFR3/MIB-1) and EORTC risk scores are predictive in primary non-muscle-invasive bladder cancer. *Eur Urol* 2010 Sep;58(3):433-41.  
<http://www.ncbi.nlm.nih.gov/pubmed/20646825>
115. Fernandez-Gomez J, Madero R, Solsona E, et al; Club Urológico Español de Tratamiento Oncológico. The EORTC tables overestimate the risk of recurrence and progression in patients with non-muscle-invasive bladder cancer treated with bacillus Calmette-Guérin: external validation of the EORTC risk tables. *Eur Urol* 2011 Sep;60(3):423-30.  
<http://www.ncbi.nlm.nih.gov/pubmed/21621906>
116. Palou J, Rodríguez-Rubio F, Millán F, et al. Recurrence at three months and high-grade recurrence as prognostic factor of progression in multivariate analysis of T1G2 bladder tumors. *Urology* 2009 Jun;73(6):1313-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/19362341>
117. Alkhateeb SS, Neill M, Bar-Moshe S, et al. Long-term prognostic value of the combination of EORTC risk group calculator and molecular markers in non-muscle-invasive bladder cancer patients treated with intravesical Bacille Calmette-Guérin. *Urol Ann* 2011 Sep;3(3):119-26.  
<http://www.ncbi.nlm.nih.gov/pubmed/21976923>
118. Lamm DL. Carcinoma in situ. *Urol Clin North Am* 1992 Aug;19(3):499-508.  
<http://www.ncbi.nlm.nih.gov/pubmed/1636234>
119. Losa A, Hurle R, Lembo A. Low dose bacillus Calmette-Guerin for carcinoma in situ of the bladder: long-term results. *J Urol* 2000 Jan;163(1):68-72.  
<http://www.ncbi.nlm.nih.gov/pubmed/10604316>
120. Griffiths TRL, Charlton M, Neal DE, et al. Treatment of carcinoma in situ with intravesical bacillus Calmette-Guerin without maintenance. *J Urol* 2002 Jun;167(6):2408-12.  
<http://www.ncbi.nlm.nih.gov/pubmed/11992047>
121. Takenaka A, Yamada Y, Miyake H, et al. Clinical outcomes of bacillus Calmette-Guérin instillation therapy for carcinoma in situ of urinary bladder. *Int J Urol* 2008 Apr;15(4):309-13.  
<http://www.ncbi.nlm.nih.gov/pubmed/18380817>

122. Solsona E, Iborra I, Dumont R, et al. The 3-month clinical response to intravesical therapy as a predictive factor for progression in patients with high risk superficial bladder cancer. *J Urol* 2000 Sep;164(3 Pt 1):685-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/10953125>
123. Van Gils-Gielen RJ, Witjes WP, Caris CT, et al. Risk factors in carcinoma in situ of the urinary bladder. *Urology* 1995 Apr;45(4):581-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/7716838>
124. Hudson MA, Herr HW. Carcinoma in situ of the bladder. *J Urol* 1995 Mar;153(3 Pt 1):564-72.  
<http://www.ncbi.nlm.nih.gov/pubmed/7861485>
125. Brausi M, Witjes JA, Lamm D, et al. A review of current guidelines and best practice recommendations for the management of nonmuscle invasive bladder cancer by the International Bladder Cancer Group. *J Urol* 2011 Dec;186(6):2158-67.  
<http://www.ncbi.nlm.nih.gov/pubmed/22014799>
126. Soloway MS, Masters S. Urothelial susceptibility to tumor cell implantation: influence of cauterization. *Cancer* 1980 Sep 1;46(5):1158-63.  
<http://www.ncbi.nlm.nih.gov/pubmed/7214299>
127. Pan JS, Slocum HK, Rustum YM, et al. Inhibition of implantation of murine bladder tumor by thiotepa in cauterized bladder. *J Urol* 1989 Dec;142(6):1589-93.  
<http://www.ncbi.nlm.nih.gov/pubmed/2511340>
128. Brocks CP, Büttner H, Böhle A. Inhibition of tumor implantation by intravesical gemcitabine in a murine model of superficial bladder cancer. *J Urol* 2005 Sep;174(3):1115-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/16094076>
129. Oosterlinck W, Kurth KH, Schröder F, et al. A prospective European Organization for Research and Treatment of Cancer Genitourinary Group randomized trial comparing transurethral resection followed by a single intravesical instillation of epirubicin or water in single stage Ta, T1 papillary carcinoma of the bladder. *J Urol* 1993 Apr;149(4):749-52.  
<http://www.ncbi.nlm.nih.gov/pubmed/8455236>
130. Sylvester RJ, Oosterlinck W, van der Meijden AP. A single immediate postoperative instillation of chemotherapy decreases the risk of recurrence in patients with stage Ta T1 bladder cancer: a metaanalysis of published results of randomized clinical trials. *J Urol* 2004 Jun;171(6 Pt 1):2186-90.  
<http://www.ncbi.nlm.nih.gov/pubmed/15126782>
131. Berrum-Svennung I, Granfors T, Jahnson S, et al. A Single instillation of epirubicin after transurethral resection of bladder tumors prevents only small recurrences. *J Urol* 2008 Jan;179(1):101-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/17997459>
132. Gudjonsson S, Adell L, Merdasa F, et al. Should all patients with non-muscle-invasive bladder cancer receive early intravesical chemotherapy after transurethral resection? The results of a prospective randomised multicentre study. *Eur Urol* 2009 Apr;55(4):773-80.  
<http://www.ncbi.nlm.nih.gov/pubmed/19153001>
133. Bouffieux C, Kurth KH, Bono A, et al. Intravesical adjuvant chemotherapy for superficial transitional cell bladder carcinoma: results of 2 European Organization for Research and Treatment of Cancer randomized trials with mitomycin C and doxorubicin comparing early versus delayed instillations and short-term versus long-term treatment. European Organization for Research and Treatment of Cancer Genitourinary Group. *J Urol* 1995 Mar;153(3 Pt 2):934-41.  
<http://www.ncbi.nlm.nih.gov/pubmed/7853578>
134. Kaasinen E, Rintala E, Hellstrom P, et al; FinnBladder Group. Factors explaining recurrence in patients undergoing chemo-immunotherapy regimens for frequently recurring superficial bladder carcinoma. *Eur Urol* 2002 Aug;42(2):167-74.  
<http://www.ncbi.nlm.nih.gov/pubmed/12160589>
135. Tolley DA, Parmar MK, Grigor KM, et al. The effect of intravesical mitomycin C on recurrence of newly diagnosed superficial bladder cancer: a further report with 7 years of follow up. *J Urol* 1996 Apr;155(4):1233-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/8632538>
136. Pode D, Alon Y, Horowitz AT, et al. The mechanism of human bladder tumor implantation in an in vitro model. *J Urol* 1986 Aug;136(2):482-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/3525861>
137. Günther JH, Jurczok A, Wulf T, et al. Optimizing syngeneic orthotopic murine bladder cancer (MB49). *Cancer Res* 1999 Jun;59(12):2834-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/10383142>

138. Böhle A, Jurczok A, Ardelt PU, et al. Inhibition of bladder carcinoma cell adhesion by oligopeptide combinations in vitro and in vivo. *J Urol* 2002 Jan;167(1):357-63.  
<http://www.ncbi.nlm.nih.gov/pubmed/11743356>
139. Oddens JR, van der Meijden AP, Sylvester R. One immediate postoperative instillation of chemotherapy in low risk Ta, T1 bladder cancer patients. Is it always safe? *Eur Urol* 2004 Sep;46(3):336-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/15306104>
140. Huncharek M, McGarry R, Kupelnick B. Impact of intravesical chemotherapy on recurrence rate of recurrent superficial transitional cell carcinoma of the bladder: results of a meta-analysis. *Anticancer Res* 2001 Jan-Feb;21(1B):765-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/11299841>
141. Böhle A, Bock PR. Intravesical bacillus Calmette-Guerin versus mitomycin C in superficial bladder cancer: formal meta-analysis of comparative studies on tumour progression. *Urology* 2004 Apr;63(4):682-6. Discussion 686-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/15072879>
142. Sylvester RJ, van der Meijden AP, Lamm DL. Intravesical bacillus Calmette-Guerin reduces the risk of progression in patients with superficial bladder cancer: a meta-analysis of the published results of randomized clinical trials. *J Urol* 2002 Nov;168(5):1964-70.  
<http://www.ncbi.nlm.nih.gov/pubmed/12394686>
143. Malmström P-U, Sylvester RJ, Crawford DE, et al. An individual patient data meta-analysis of the long-term outcome of randomised studies comparing intravesical mitomycin C versus bacillus Calmette-Guérin for non-muscle-invasive bladder cancer. *Eur Urol* 2009 Aug;56(2):247-56.  
<http://www.ncbi.nlm.nih.gov/pubmed/19409692>
144. Sylvester RJ, Brausi MA, Kirkels WJ, et al; EORTC Genito-Urinary Tract Cancer Group. Long-term efficacy results of EORTC genito-urinary group randomized phase 3 study 30911 comparing intravesical instillations of epirubicin, bacillus Calmette-Guérin, and bacillus Calmette-Guérin plus isoniazid in patients with intermediate- and high-risk stage Ta T1 urothelial carcinoma of the bladder. *Eur Urol* 2010 May;57(5):766-73  
<http://www.ncbi.nlm.nih.gov/pubmed/20034729>
145. Shang PF, Kwong J, Wang ZP, et al. Intravesical Bacillus Calmette-Guérin versus epirubicin for Ta and T1 bladder cancer. *Cochrane Database Syst Rev* 2011 May 11;(5):CD006885.  
<http://www.ncbi.nlm.nih.gov/pubmed/21563157>
146. Sylvester RJ, Oosterlinck W, Witjes JA. The schedule and duration of intravesical chemotherapy in patients with non muscle invasive bladder cancer: a systematic review of the published results of randomized clinical trials. *Eur Urol* 2008 Apr;53(4):709-19.  
<http://www.ncbi.nlm.nih.gov/pubmed/18207317>
147. Lammers RJ, Witjes JA, Inman BA, et al. The role of a combined regimen with intravesical chemotherapy and hyperthermia in the management of non-muscle-invasive bladder cancer: a systematic review. *Eur Urol* 2011 Jul;60(1):81-93.  
<http://www.ncbi.nlm.nih.gov/pubmed/21531502>
148. Di Stasi SM, Giannantoni A, Giurioli A, et al. Sequential BCG and electromotive mitomycin versus BCG alone for high-risk superficial bladder cancer: a randomised controlled trial. *Lancet Oncol* 2006 Jan;7(1):43-51.  
<http://www.ncbi.nlm.nih.gov/pubmed/16389183>
149. Au JL, Baladament RA, Wientjes MG, et al; International Mitomycin C Consortium. International Mitomycin C Consortium. Methods to improve efficacy of intravesical mitomycin C: results of a randomized phase III trial. *J Natl Cancer Inst* 2001 Apr;93(8):597-604.  
<http://www.ncbi.nlm.nih.gov/pubmed/11309436>
150. Giesbers AA, Van Helsdingen PJ, Kramer AE. Recurrence of superficial bladder carcinoma after intravesical instillation of mitomycin-C. Comparison of exposure times. *Br J Urol* 1989 Feb;63(2):176-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/2495144>
151. Kuroda M, Niijima T, Kotake T, et al; 6th Trial of the Japanese Urological Cancer Research Group. Effect of prophylactic treatment with intravesical epirubicin on recurrence of superficial bladder cancer -The 6th Trial of the Japanese Urological Cancer Research Group (JUICRG): a randomized trial of intravesical epirubicin at dose of 20mg/40ml, 30 mg/40 ml, 40 mg/40 ml. *Eur Urol* 2004 May;45(5):600-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/15082202>

152. Shelley MD, Kynaston H, Court J, et al. A systematic review of intravesical bacillus Calmette-Guérin plus transurethral resection vs transurethral resection alone in Ta and T1 bladder cancer. *BJU Int* 2001 Aug;88(3):209-16.  
<http://www.ncbi.nlm.nih.gov/pubmed/11488731>
153. Han RF, Pan JG. Can intravesical bacillus Calmette-Guérin reduce recurrence in patients with superficial bladder cancer? A meta-analysis of randomized trials. *Urology* 2006 Jun;67(6):1216-23.  
<http://www.ncbi.nlm.nih.gov/pubmed/16765182>
154. Shelley MD, Wilt TJ, Court J, et al. Intravesical bacillus Calmette-Guérin is superior to mitomycin C in reducing tumour recurrence in high-risk superficial bladder cancer: a metaanalysis of randomized trials. *BJU Int* 2004 Mar;93(4):485-90.  
<http://www.ncbi.nlm.nih.gov/pubmed/15008714>
155. Böhle A, Jocham D, Bock PR. Intravesical bacillus Calmette-Guerin versus mitomycin C for superficial bladder cancer: a formal meta-analysis of comparative studies on recurrence and toxicity. *J Urol* 2003 Jan;169(1):90-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/12478111>
156. Duchek M, Johansson R, Jahnson S, et al. Bacillus Calmette-Guérin Is superior to a combination of epirubicin and interferon- $\alpha$ 2b in the intravesical treatment of patients with stage T1 urinary bladder cancer. a prospective, randomized, nordic study. *Eur Urol* 2010 Jan;57(1):25-31.  
<http://www.ncbi.nlm.nih.gov/pubmed/19819617>
157. Järvinen R, Kaasinen E, Sankila A, et al. Long-term efficacy of maintenance bacillus Calmette-Guérin versus maintenance mitomycin C instillation therapy in frequently recurrent TaT1 tumours without carcinoma in situ: a subgroup analysis of the prospective, randomised FinnBladder I study with a 20-year follow-up. *Eur Urol* 2009 Aug;56(2):260-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/19395154>
158. Huncharek M, Kupelnick B. The influence of intravesical therapy on progression of superficial transitional cell carcinoma of the bladder: a metaanalytic comparison of chemotherapy versus bacilli Calmette-Guerin immunotherapy. *Am J Clin Oncol* 2004 Oct;27(5):522-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/15596924>
159. Oddens J, Brausi M, Sylvester R, et al. Final Results of an EORTC-GU Cancers Group Randomized Study of Maintenance Bacillus Calmette-Guérin in Intermediate- and High-risk Ta, T1 Papillary Carcinoma of the Urinary Bladder: One-third Dose Versus Full Dose and 1 Year Versus 3 Years of Maintenance. *Eur Urol* 2013 Mar;63(3):462-72.  
<http://www.ncbi.nlm.nih.gov/pubmed/23141049>
160. Morales, A, Eiding D, Bruce AW. Intracavitary bacillus Calmette-Guerin in the treatment of superficial bladder tumors. *J Urol* 1976 Aug;116(2):180-3.  
<http://www.ncbi.nlm.nih.gov/pubmed/820877>
161. Lamm DL, Blumenstein BA, Crissman JD, et al. Maintenance bacillus Calmette-Guerin immunotherapy for recurrent Ta, T1 and carcinoma in situ transitional cell carcinoma of the bladder: a randomized Southwest Oncology Group Study. *J Urol* 2000 Apr;163(4):1124-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/10737480>
162. Zlotta AR, van Vooren JP, Huygen K, et al. What is the optimal regimen for BCG intravesical therapy? Are six weekly instillations necessary? *Eur Urol* 2000 Apr;37(4):470-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/10765079>
163. Van der Meijden AP, Sylvester RJ, Oosterlinck W, et al; EORTC Genito-Urinary Tract Cancer Group. Maintenance bacillus Calmette-Guerin for Ta, T1 bladder tumours is not associated with increased toxicity: results from a European Organisation for Research and Treatment of Cancer Genito-Urinary Group Phase III Trial. *Eur Urol* 2003 Oct;44(4):429-34.  
<http://www.ncbi.nlm.nih.gov/pubmed/14499676>
164. Herr HW. Intravesical bacillus Calmette-Guérin outcomes in patients with bladder cancer and asymptomatic bacteriuria. *J Urol* 2012 Feb;187(2):435-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/22177154>
165. Herr HW. Outpatient urological procedures in antibiotic-naive patients with bladder cancer with asymptomatic bacteriuria. *BJU Int* 2012 Aug 9.  
<http://www.ncbi.nlm.nih.gov/pubmed/22883017>
166. Lamm DL, van der Meijden PM, Morales A, et al. Incidence and treatment of complications of bacillus Calmette-Guerin intravesical therapy in superficial bladder cancer. *J Urol* 1992 Mar;147(3):596-600.  
<http://www.ncbi.nlm.nih.gov/pubmed/1538436>
167. Palou J, Angerri O, Segarra J, et al. Intravesical bacillus Calmette-Guèrin for the treatment of superficial bladder cancer in renal transplant patients. *Transplantation* 2003 Nov 27;76(10):1514-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/14657696>

168. Yossepowitch O, Eggener SE, Bochner BH, et al. Safety and efficacy of intravesical bacillus Calmette-Guerin instillations in steroid treated and immunocompromised patients. *J Urol* 2006 Aug;176(2):482-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/16813873>
169. Rodríguez F, Palou J, Martínez R, et al. Practical guideline for the management of adverse events associated with BCG installations. *Arch Esp Urol*. 2008 Jun;61(5):591-6. (Article in Spanish)  
<http://www.ncbi.nlm.nih.gov/pubmed/18709813>
170. Witjes JA, Palou J, Soloway M, et al. Clinical practice recommendations for the prevention and management of intravesical therapy-associated adverse events. *Eur Urol Suppl* 2008 Oct;7(10):667-74.
171. Tinazzi E, Ficarra V, Simeoni S, et al. Reactive arthritis following BCG immunotherapy for urinary bladder carcinoma: a systematic review. *Rheumatol Int* 2006 Apr;26(6):481-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/16220289>
172. Falkensammer C, Gozzi C, Hager M, et al. Late occurrence of bilateral tuberculous-like epididymo-orchitis after intravesical bacille Calmette-Guerin therapy for superficial bladder carcinoma. *Urology* 2005 Jan;65(1):175.  
<http://www.ncbi.nlm.nih.gov/pubmed/15667898>
173. Palou J, Rodríguez-Villamil L, Andreu-Crespo A, et al., Intravesical treatment of severe bacillus Calmette-Guerin cystitis. *Int Urol Nephrol* 2001;33(3):485-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/12230277>
174. Martínez-Pineiro JA, Flores N, Isorna S, et al. Long-term follow-up of a randomized prospective trial comparing a standard 81 mg dose of intravesical bacille Calmette-Guerin with a reduced dose of 27 mg in superficial bladder cancer. *BJU Int* 2002 May;89(7):671-80.  
<http://www.ncbi.nlm.nih.gov/pubmed/11966623>
175. Martínez-Piñero JA, Martínez-Piñero L, Solsona E, et al. Has a 3-fold decreased dose of bacillus Calmette-Guerin the same efficacy against recurrences and progression of T1G3 and Tis bladder tumors than the standard dose? Results of a prospective randomized trial. *J Urol* 2005 Oct;174(4 Pt 1):1242-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/16145378>
176. Ojea A, Nogueira JL, Solsona E, et al; CUETO Group (Club Urológico Español De Tratamiento Oncológico). A multicentre, randomised prospective trial comparing three intravesical adjuvant therapies for intermediate-risk superficial bladder cancer: low-dose bacillus Calmette-Guerin (27mg) versus very low-dose bacillus Calmette-Guerin (13.5mg) versus mitomycin C. *Eur Urol* 2007 Nov;52(5):1398-406.  
<http://www.ncbi.nlm.nih.gov/pubmed/17485161>
177. Van der Meijden APM, Sylvester R, Oosterlinck W, et al; for the EAU Working Party on Non Muscle Invasive Bladder Cancer. EAU guidelines on the diagnosis and treatment of urothelial carcinoma in situ. *Eur Urol* 2005 Sep;48(3):363-71.  
<http://www.ncbi.nlm.nih.gov/pubmed/15994003>
178. Jakse G, Hall R, Bono A, et al and members of the EORTC GU Group. Intravesical BCG in patients with carcinoma in situ of the urinary bladder: long-term results of EORTC GU Group phase II protocol 30861. *Eur Urol* 2001 Aug;40(2):144-50.  
<http://www.ncbi.nlm.nih.gov/pubmed/11528191>
179. Gofrit ON, Pode D, Pizov G, et al. The natural history of bladder carcinoma in situ after initial response to bacillus Calmette-Guérin immunotherapy. *Urol Oncol* 2009 May-Jun;27(3):258-62.  
<http://www.ncbi.nlm.nih.gov/pubmed/18440839>
180. Sylvester RJ, van der Meijden APM, Witjes JA, et al. Bacillus Calmette-Guerin versus chemotherapy in the intravesical treatment of patients with carcinoma in situ of the bladder: a meta-analysis of the published results of randomized clinical trials. *J Urol* 2005 Jul;174(1):86-92.  
<http://www.ncbi.nlm.nih.gov/pubmed/15947584>
181. Kaasinen E, Wijkstrom H, Malmstrom PU, et al. Alternating mitomycin C and BCG instillations versus BCG alone in treatment of carcinoma in situ of the urinary bladder: a Nordic study. *Eur Urol* 2003 Jun;43(6):637-45.  
<http://www.ncbi.nlm.nih.gov/pubmed/12767365>
182. Solsona E, Iborra I, Ricos JV, et al. Extravesical involvement in patients with bladder carcinoma in situ: biological and therapy implications. *J Urol* 1996 Mar;155(3):895-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/8583601>
183. Palou J, Baniel J, Klotz L, et al. Urothelial carcinoma of the prostate. *Urology* 2007 Jan;69(1 Suppl):50-61.  
<http://www.ncbi.nlm.nih.gov/pubmed/17280908>

184. Palou Redorta J, Schatteman P, Huguet Pérez J, et al. Intravesical instillations with bacillus calmette-guérin for the treatment of carcinoma in situ involving prostatic ducts. *Eur Urol* 2006 May;49(5):834-8; discussion 838.  
<http://www.ncbi.nlm.nih.gov/pubmed/16426729>
185. Herr HW, Dalbagni G. Defining bacillus Calmette-Guerin refractory superficial bladder tumours. *J Urol* 2003 May;169(5):1706-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/12686813>
186. Lerner SP, Tangen CM, Sucharew H, et al. Failure to achieve a complete response to induction BCG therapy is associated with increased risk of disease worsening and death in patients with high risk non muscle invasive bladder cancer. *Urol Oncol* 2009 Mar-Apr;27(2):155-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/18367117>
187. van den Bosch S, Alfred Witjes J. Long-term cancer-specific survival in patients with high-risk, non-muscle-invasive bladder cancer and tumour progression: a systematic review. *Eur Urol* 2011 Sep;60(3):493-500.  
<http://www.ncbi.nlm.nih.gov/pubmed/21664041>
188. Gallagher BL, Joudi FN, Maymi JL, et al. Impact of previous bacille Calmette-Guérin failure pattern on subsequent response to bacille Calmette-Guérin plus interferon intravesical therapy. *Urology* 2008 Feb;71(2):297-301.  
<http://www.ncbi.nlm.nih.gov/pubmed/18308107>
189. Rosevear HM, Lightfoot AJ, Birusingh KK, et al. Factors affecting response to bacillus Calmette-Guérin plus interferon for urothelial carcinoma in situ. *J Urol* 2011 Sep;186(3):817-23.  
<http://www.ncbi.nlm.nih.gov/pubmed/21788050>
190. Yates DR, Brausi MA, Catto JW, et al. Treatment options available for bacillus Calmette-Guérin failure in non-muscle-invasive bladder cancer. *Eur Urol* 2012 Dec;62(6):1088-96.  
<http://www.ncbi.nlm.nih.gov/pubmed/22959049>
191. Dalbagni G, Russo P, Bochner B, et al. Phase II trial of intravesical gemcitabine in bacille Calmette-Guerin-refractory transitional cell carcinoma of the bladder. *J Clin Oncol* 2006 Jun;24(18):2729-34.  
<http://www.ncbi.nlm.nih.gov/pubmed/16782913>
192. Mohanty NK, Nayak RL, Vasudeva P, et al. Intravesicle gemcitabine in management of BCG refractory superficial TCC of urinary bladder-our experience. *Urol Oncol* 2008 Nov-Dec;26(6):616-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/18367121>
193. Barlow L, McKiernan J, Sawczuk I, et al. A single-institution experience with induction and maintenance intravesical docetaxel in the management of non-muscle-invasive bladder cancer refractory to bacilli Calmette-Guerin therapy. *BJU Int* 2009 Oct;104(8):1098-102.  
<http://www.ncbi.nlm.nih.gov/pubmed/19389012>
194. Steinberg G, Bahnson R, Brosman S, et al. Efficacy and safety of valrubicin for the treatment of bacillus Calmette-Guerin refractory carcinoma in situ of the bladder. The Valrubicin Study Group. *J Urol* 2000 Mar;163(3):761-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/10687972>
195. Nativ O, Witjes JA, Hendricksen K, et al. Combined thermo-chemotherapy for recurrent bladder cancer after bacillus Calmette-Guerin. *J Urol* 2009 Oct; 182(4):1313-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/19683278>
196. Joudi FN, Smith BJ, O'Donnell MA. Final results from a national multicenter phase II trial of combination bacillus Calmette-Guerin plus interferon alpha-2B for reducing recurrence of superficial bladder cancer. *Urol Oncol* 2006 Jul-Aug;24(4):344-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/16818189>
197. Di Lorenzo G, Perdonà S, Damiano R, et al. Gemcitabine versus bacille Calmette-Guérin after initial bacille Calmette-Guérin failure in non-muscle-invasive bladder cancer: a multicenter prospective randomized trial. *Cancer* 2010 Apr;116(8):1893-900.  
<http://www.ncbi.nlm.nih.gov/pubmed/20162706>
198. Yates DR, Rouprêt M. Failure of bacille Calmette-Guérin in patients with high risk non-muscle-invasive bladder cancer unsuitable for radical cystectomy: an update of available treatment options. *BJU International* 2010 Jul;106(2):162-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/20201829>
199. Zlotta AR, Fleshner NE, Jewett MA. The management of BCG failure in non-muscle-invasive bladder cancer: an update. *Can Urol Assoc J* 2009 Dec;3(6 Suppl 4):S199-205.  
<http://www.ncbi.nlm.nih.gov/pubmed/20019985>

200. Turker P, Bostrom PJ, Wroclawski ML, et al. Upstaging of urothelial cancer at the time of radical cystectomy: factors associated with upstaging and its effect on outcome. *BJU Int* 2012 Sep;110(6):804-11.  
<http://www.ncbi.nlm.nih.gov/pubmed/22321341>
201. Chalasani V, Kassouf W, Chin JL, et al. Radical cystectomy for the treatment of T1 bladder cancer: the Canadian Bladder Cancer Network experience. *Can Urol Assoc J.* 2011 Apr;5(2):83-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/21470529>
202. May M, Bastian PJ, Brookman-May S, et al. Pathological upstaging detected in radical cystectomy procedures is associated with a significantly worse tumour-specific survival rate for patients with clinical T1 urothelial carcinoma of the urinary bladder. *Scand J Urol Nephrol* 2011 Sep;45(4):251-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/21388337>
203. Svatek RS, Shariat SF, Novara G, et al. Discrepancy between clinical and pathological stage: external validation of the impact on prognosis in an international radical cystectomy cohort. *BJU Int* 2011 Mar;107(6):898-904.  
<http://www.ncbi.nlm.nih.gov/pubmed/21244604>
204. Gupta A, Lotan Y, Bastian PJ, et al. Outcomes of patients with clinical T1 grade 3 urothelial cell bladder carcinoma treated with radical cystectomy. *Urology* 2008 Feb;71(2):302-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/18308108>
205. Denzinger S, Fritsche HM, Otto W, et al. Early versus deferred cystectomy for initial high-risk pT1G3 urothelial carcinoma of the bladder: do risk factors define feasibility of bladder-sparing approach? *Eur Urol* 2008 Jan;53(1):146-52.  
<http://www.ncbi.nlm.nih.gov/pubmed/17624657>
206. Lambert EH, Pierorazio PM, Olsson CA, et al. The increasing use of intravesical therapies for stage T1 bladder cancer coincides with decreasing survival after cystectomy. *BJU Int* 2007 Jul;100(1):33-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/17552951>
207. Shariat SF, Palapattu GS, Karakiewicz PI, et al. Discrepancy between clinical and pathologic stage: impact on prognosis after radical cystectomy. *Eur Urol* 2007 Jan;51(1):137-49; discussion 149-51.  
<http://www.ncbi.nlm.nih.gov/pubmed/16793197>
208. Thalmann GN, Markwalder R, Shahin O, et al. Primary T1G3 bladder cancer: organ preserving approach or immediate cystectomy? *J Urol* 2004 Jul;172(1):70-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/15201740>
209. Amling CL, Thrasher JB, Frazier HA, et al. Radical cystectomy for stages Ta, Tis and T1 transitional cell carcinoma of the bladder. *J Urol* 1994 Jan;151(1):31-5; discussion 35-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/8254828>
210. Freeman JA, Esrig D, Stein JP, et al. Radical cystectomy for high risk patients with superficial bladder cancer in the era of orthotopic urinary reconstruction. *Cancer* 1995 Sep 1;76(5):833-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/8625187>
211. Bianco FJ Jr, Justa D, Grignon DJ, et al. Management of clinical T1 bladder transitional cell carcinoma by radical cystectomy. *Urol Oncol* 2004 Jul-Aug;22(4):290-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/15283885>
212. Herr HW, Sogani PC. Does early cystectomy improve the survival of patients with high risk superficial bladder tumors? *J Urol* 2001 Oct;166(4):1296-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/11547061>
213. Raj GV, Herr H, Serio AM, et al. Treatment paradigm shift may improve survival of patients with high risk superficial bladder cancer. *J Urol* 2007 Apr;177(4):1283-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/17382713>
214. Stein JP, Lieskovsky G, Cote R, et al. Radical cystectomy in the treatment of invasive bladder cancer: long-term results in 1,054 patients. *J Clin Oncol* 2001 Feb;19(3):666-75.  
<http://www.ncbi.nlm.nih.gov/pubmed/11157016>
215. Hautmann RE, Gschwend JE, de Petriconi RC, et al. Cystectomy for transitional cell carcinoma of the bladder: results of surgery only series in the neobladder era. *J Urol* 2006 Aug;176(2):486-92.  
<http://www.ncbi.nlm.nih.gov/pubmed/16813874>
216. Ghoneim MA, Abdel-Latif M, El-Mekresh M, et al, Radical cystectomy for carcinoma of the bladder: 2,720 consecutive cases 5 year later. *J Urol* 2008 Jul;180(1):121-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/18485392>
217. Madersbacher S, Hochreiter W, Burkhard F, et al. Radical cystectomy for bladder cancer today - a homogeneous series without neoadjuvant therapy. *J Clin Oncol* 2003 Feb;21(4):690-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/12586807>



218. Shariat SF, Karakiewicz PI, Palapattu GS, et al. Outcomes of radical cystectomy for transitional cell carcinoma of the bladder: a contemporary series from the bladder cancer research consortium. *J Urol* 2006 Dec;176(6 Pt 1):2414-22.  
<http://www.ncbi.nlm.nih.gov/pubmed/17085118>
219. Shariat SF, Palapattu GS, Amiel GE, et al. Characteristics and outcomes of patients with carcinoma in situ only at radical cystectomy. *Urology* 2006 Sep;68(3):538-42.  
<http://www.ncbi.nlm.nih.gov/pubmed/16979748>
220. Oge O, Erdem E, Atsu N, et al. Proposal for changes in cystoscopic follow-up of patients with low-grade pTa bladder tumour. *Eur Urol* 2000 Mar;37(3):271-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/10720851>
221. Holmäng S, Andius P, Hedelin H, et al. Stage progression in Ta papillary urothelial tumours: relationship to grade, immunohistochemical expression of tumour markers, mitotic frequency and DNA ploidy. *J Urol* 2001 Apr;165(4):1124-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/11257652>
222. Fujii Y, Kawakami S, Koga F, et al. Long-term outcome of bladder papillary urothelial neoplasms of low malignant potential. *BJU Int* 2003 Oct;92(6):559-62.  
<http://www.ncbi.nlm.nih.gov/pubmed/14511033>
223. Leblanc B, Duclos AJ, Bénard F, et al. Long-term follow-up of initial Ta grade 1 transitional cell carcinoma of the bladder. *J Urol* 1999 Dec;162(6):1946-50.  
<http://www.ncbi.nlm.nih.gov/pubmed/10569544>
224. Zieger K, Wolf H, Olsen PR, et al. Long-term follow-up of non-invasive bladder tumours (stage Ta): recurrence and progression. *BJU Int* 2000 May;85(7):824-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/10792160>
225. Borhan A, Reeder JE, O'Connell MJ, et al. Grade progression and regression in recurrent urothelial cancer. *J Urol* 2003 Jun;169(6):2106-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/12771728>
226. Soloway M, Bruck DS, Kim SS. Expectant management of small recurrent, non-invasive papillary bladder tumours. *J Urol* 2003 Aug;170(2 Pt 1):438-41.  
<http://www.ncbi.nlm.nih.gov/pubmed/12853794>
227. Gofrit ON, Pode D, Lazar A, et al. Watchful waiting policy in recurrent Ta G1 bladder tumours. *Eur Urol* 2006 Feb;49(2):303-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/16413659>
228. Pruthi RS, Baldwin N, Bhalani V, et al. Conservative Management of Low Risk Superficial Bladder Tumors. *J Urol* 2008 Jan;179(1):87-90.  
<http://www.ncbi.nlm.nih.gov/pubmed/17997444>
229. Holmang S, Johansson SL. Stage Ta-T1 bladder cancer: the relationship between findings at first follow-up cystoscopy and subsequent recurrence and progression. *J Urol* 2002 Apr;167(4):1634-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/11912378>
230. Mariappan P, Smith G. A surveillance schedule for G1Ta bladder cancer allowing efficient use of check cystoscopy and safe discharge at 5 years based on a 25-year prospective database. *J Urol* 2005 Apr;173(4):1008-11.  
<http://www.ncbi.nlm.nih.gov/pubmed/15758711>
231. Soukup V, Babjuk M, Bellmunt J, et al. Follow-up after surgical treatment of bladder cancer: a critical analysis of the literature. *Eur Urol* 2012 Aug;62(2):290-302.  
<http://www.ncbi.nlm.nih.gov/pubmed/22609313>
232. Holmäng S, Ströck V. Should follow-up cystoscopy in bacillus Calmette-Guérin-treated patients continue after five tumour-free years? *Eur Urol* 2012 Mar;61(3):503-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/22119022>

## 11. ABBREVIATIONS USED IN THE TEXT

*This list is not comprehensive for the most common abbreviations.*

5-ALA	5-aminolaevulinic acid
ASR	age standardised incidence rate
BCG	bacillus Calmette-Guérin
BTA	bladder tumour antigen
CIS	carcinoma <i>in situ</i>
CT	computed tomography
CUETO Club	Urológico Español de Tratamiento Oncológico (Spanish Oncology Group)
EAU	European Association of Urology
EORTC	European Organization for Research and Treatment of Cancer
EORTC-GUCG	EORTC Genito-Urinary Cancer Group
FISH	fluorescence in situ hybridisation
GR	grade of recommendation
HAL	hexaminolaevulinic acid
ISUP	International Society of Urological Pathology
IVU	intravenous urography
LE	level of evidence
MMC	mitomycin C
NMIBC	non-muscle-invasive bladder cancer
NVP	negative predictive value
PDD	photodynamic diagnosis
PUNLMP	papillary urothelial neoplasms of low malignant potential
RCT	randomised controlled trial
TCC	transitional cell carcinoma
TNM	tumour, node, metastasis
TUR	transurethral resection
UICC	Union International Contre le Cancer
US	ultrasonography
WHO	World Health Organization

### **Conflict of interest**

All members of the Non-Muscle-Invasive Bladder Cancer guidelines working group have provided disclosure statements of all relationships that they have that might be perceived as a potential source of a conflict of interest. This information is publically accessible through the European Association of Urology website. This guidelines document was developed with the financial support of the European Association of Urology. No external sources of funding and support have been involved. The EAU is a non-profit organisation and funding is limited to administrative assistance and travel and meeting expenses. No honoraria or other reimbursements have been provided.